NYSDOH Adult Sepsis and COVID-19 Data Dictionary

Digitalized Data Collection, D2.1

Version (Digital) D2.1

<mark>March</mark> 4,</mark> 2021

This dictionary includes the administrative codes found in the Appendices in a CSV format available for download to assist in data extraction.

The most recent version of this document, the *Frequently Asked Questions* document, the *Table of Elements* data template, and the instructions may be found at: <u>https://ny.sepsis.ipro.org</u>

Questions regarding this document should be submitted at: <u>https://ny.sepsis.ipro.org/support</u>

Prepared by: Office of Quality and Patient Safety New York State Department of Health Released for use 2020 Revised March 4, 2021

Table of Contents

Key points to remember during data extraction	
Inclusion Definition	10
Demographic Variables	13
Admission Datetime	14
Arrival Datetime	15
Date of Birth	
Discharge Datetime	
Discharge Status	19
Ethnicity	21
Facility Identifier	22
Gender	23
ICD-10-CM Code (n)	24
ICD-10-CM POA Indicator (n)	25
Insurance Number	26
Medical Record Number	27
Other Payer	
Patient Control Number	29
Patient Zip Code of Residence	
Payer	
Race	
Source of Admission	
Transferred In	
Transferred Out	35
Transfer Facility Identifier Receiving	
Transfer Facility Identifier Sending	
Transfer Facility Name Receiving	
Transfer Facility Name Sending	
Unique Personal Identifier	40
Comorbidity/Risk Factor (POA) Variables	41
Acute Cardiovascular Conditions	42

AIDS/HIV Disease	43
Altered Mental Status	44
Asthma	45
Chronic Liver Disease	46
Chronic Renal Failure	47
Chronic Respiratory Failure	48
Coagulopathy	49
Congestive Heart Failure	50
COPD	51
Dementia	52
Diabetes	53
Dialysis Comorbidity	54
History of COVID -19	55
History of COVID-19 Datetime	56
History of Other Cardiovascular Disease	57
Hypertension	58
Immunocompromising	59
Lymphoma Leukemia Multiple Myeloma	60
Mechanical Ventilation Comorbidity	61
Medication Anticoagulation	62
Medication Immune Modifying	63
Metastatic Cancer	64
Obesity	65
Patient Care Considerations	66
Patient Care Considerations Date	67
Pregnancy Comorbidity	68
Pregnancy Status During Hospitalization	69
Smoking Vaping	70
Tracheostomy on Arrival	71
Clinical Variables	72
COVID-19 Exposure	73
COVID-19 Virus	74
Drug Resistant Pathogen	75

Flu Positive	76
Suspected Source of Infection	77
Treatment (during hospitalization) Variables	78
Dialysis Treatment	79
During Hospital Anticoagulation	80
During Hospital Immune Modifying Medication	81
During Hospital Remdesivir	82
ECMO	83
High Flow Nasal Cannula	84
Mechanical Ventilation Treatment	85
Non-Invasive Positive Pressure Ventilation	86
Vasopressor Administration	87
Outcome (at discharge) Variables	88
Dialysis Outcome	89
Mechanical Ventilation Outcome	90
Tracheostomy at Discharge	91
Outcome (during hospitalization) Variables	92
Cardiovascular Outcomes	93
ICU During Hospitalization	94
PE/DVT	95
Tracheostomy in Hospital	96
Severity Variables	97
aPTT 1	
aPTT 2	
aPTT 3	
aPTT Max	
aPTT Datetime 1	
aPTT Datetime 2	
aPTT Datetime 3	
aPTT Datetime Max	
Diastolic First	102
Diastolic Second	102
Diastolic Third	

Diastolic Min	
Diastolic First Datetime 1	
Diastolic Second Datetime 2	
Diastolic Third Datetime 3	
Diastolic Datetime Min	
INR 1	
INR 2	
INR 3	
INR Max	
INR Datetime 1	
INR Datetime 2	
INR Datetime 3	
INR Datetime Max	
Lactate Level 1	
Lactate Level 2	
Lactate Level 3	
Lactate Level Max	
Lactate Level Datetime 1	
Lactate Level Datetime 2	
Lactate Level Datetime 3	
Lactate Level Datetime Max	
Organ Dysfunction CNS	
Organ Dysfunction Hepatic Arrival	115
Organ Dysfunction Hepatic Max	115
Organ Dysfunction Hepatic Arrival Datetime	116
Organ Dysfunction Hepatic Max Datetime	
Organ Dysfunction Renal Arrival	
Organ Dysfunction Renal Max	
Organ Dysfunction Renal Arrival Datetime	
Organ Dysfunction Renal Max Datetime	
Organ Dysfunction Respiratory	
Platelets 1	
Platelets 2	

Platelets 3	
Platelets Min	
Platelets Datetime 1	125
Platelets Datetime 2	125
Platelets Datetime 3	125
Platelets Datetime Min	125
SIRS Heart Rate 1	127
SIRS Heart Rate 2	127
SIRS Heart Rate 3	127
SIRS Heart Rate Max	127
SIRS Heart Rate Datetime 1	129
SIRS Heart Rate Datetime 2	129
SIRS Heart Rate Datetime 3	129
SIRS Heart Rate Datetime Max	129
SIRS Leukocyte Arrival	131
SIRS Leukocyte Min	131
SIRS Leukocyte Max	131
SIRS Leukocyte Arrival Datetime	133
SIRS Leukocyte Min Datetime	133
SIRS Leukocyte Max Datetime	133
SIRS Respiratory Rate 1	135
SIRS Respiratory Rate 2	135
SIRS Respiratory Rate 3	135
SIRS Respiratory Rate Max	135
SIRS Respiratory Rate Datetime 1	137
SIRS Respiratory Rate Datetime 2	137
SIRS Respiratory Rate Datetime 3	137
SIRS Respiratory Rate Datetime Max	137
SIRS Temperature 1	139
SIRS Temperature 2	
SIRS Temperature 3	139
SIRS Temperature Max	
SIRS Temperature Datetime 1	

SIRS Temperature Datetime 2	141
SIRS Temperature Datetime 3	141
SIRS Temperature Datetime Max	141
Systolic First	143
Systolic Second	143
Systolic Third	143
Systolic Min	143
Systolic First Datetime 1	145
Systolic Second Datetime 2	145
Systolic Third Datetime 3	145
Systolic Datetime Min	145
Change Log	147

Key points to remember during data extraction

The New York State Department of Health (NYSDOH) is seeking the collection of data for all inpatient, Emergency Department (ED), and observation patients for severe sepsis, septic shock, and severe COVID-19 patients evaluated in the hospital (including ED, Observation, and inpatient patients) as defined by the case *inclusion definition* provided on the following page of this dictionary.

Data for all patients who are 21 years of age or older are to be reported into the adult NYSDOH database. Patient age at admission should be used to determine reporting to the adult or the pediatric database. If a patient is observation only or ED only, please use patient age at the time of arrival for determination of the adult or pediatric database inclusion.

When using the appendices for the identification of relevant ICD-10-CM codes, be sure to capture any code (ICD-10-CM) in any position at any point during hospitalization unless otherwise indicated in the variable directions.

Some variables are required but there are some exceptions such as *Transfer Facility Identifier Receiving* and *Transfer Facility Identifier Sending* which are situational. For transfer data elements we recognize that your hospital EHR may not have Transfer Facility Identifier Receiving/Sending but may have Transer Facility Name Receiving/Sending. Please report all data you have regarding transfers. Within hospital (interdepartment transfers) are not considered transfers for these data elements.

Hospitals that have within hospital transfers patients (i.e. patient transferred from one unit to another within the same hospital) should report the case as it is collected in the EHR. For example, if your EHR represents a patient transferred from a rehab unit to an acute care unit as one singe admission in your EHR, please report this admission as one record. If this is reported as two separate admissions, please submit it as two separate records. Be sure to use the appropariate discharge disposition to accurately represent the case.

This data dictionary has been designed to eliminate the need for manual chart abstraction and to permit hospitals to utilize their information technology staff and electronic medical record systems to extract the necessary data. This data will be accepted in to the current portal in a flat file format following existing procedures which may be found at https://ny.sepsis.ipro.org/.

CSV files of codes in appendices are provided separately. Each CSV file contains three columns: the codes of the variable/data element in appendices, the corresponding code description, and the subcategory if applicable. For example:

ICD-10-CM CODE	ICD-10-CM CODE DESCRIPTION	Subcategory
12101	ST elevation (STEMI) myocardial infarction	MI
	involving left main coronary artery	
12102	ST elevation (STEMI) myocardial infarction	MI
	involving left anterior descending coronary	
	artery	

Please note that data elements with multiple selections (more than Yes/No) will have values/contents in the subcategory column in the CSV files, for example *Acute Cardiovascular Conditions*. In general, the naming convention for CSV files is TemplateVariable_code_Version. For example:

• asthma_code_VerD2.1.csv

In the event that the CSV files are for NDC codes of medications, ndc is added in the naming convention. For example:

medication_anticoagulation_ndc_code_VerD2.1.csv

Inclusion Definition

The NYSDOH is identifying the (denominator) population of cases for inclusion into the database using ICD-10-CM codes. Hospitals may use all sources of data for case inclusion (electronic medical record codes as well as administrative and billing codes). This will allow for electronic identification of cases. The ICD-10-CM code-based definition for identifying the severe sepsis/septic shock and severe COVID-19 patient population for abstraction includes the following codes which are presented in Tables A and B. Cases with codes in either table are to be reported.

Hospitals will report cases where criteria are met by:

- At least one code in Table A alone; OR
- At least two codes in Table B, one of which must be either U071 OR U072 as well as one or more of the codes beginning with J80 through T8112XA

Examples:

- Patient with Code T8112XA and no other code from Table A or Table B is reported.
- Patient with U072 and R602 and no other code from Table A or Table B is reported.
- Patient with U071 and no other code from Table A or Table B is <u>not</u> reported.
- Patient with R6520 and no other codes from Table A or Table B is reported. This case is reported because R6520 alone is a reportable case regardless of additional codes reported on the case.

Table A: Severe sepsis and/or septic shock inclusion ICD-10-CM codes

Severe Sepsis/Septic Shock		
ICD-10-CM Description		
R6520 Severe sepsis without septic shock		
R6521 Severe sepsis with septic shock		
T8112XA	Post procedural septic shock, initial encounter	

OR

Table B: Severe COVID-19 inclusion ICD-10-CM codes

Severe COVI	D-19	
ICD-10-CM	Description	Туре
U071	COVID-19, virus identified	COVID-19
U072	COVID-19, virus not identified (Clinically-epidemiologically diagnosed COVID-19)	COVID-19
J1282	Pneumonia due to coronavirus disease 2019 (This code is effective as of January 1, 2021).	COVID-19
AND (one or more of the following)		
J80	Acute respiratory distress	Respiratory

Severe COVI	D-19	
ICD-10-CM	Description	Туре
	syndrome	
J9600	Acute respiratory failure, unsp w	Respiratory
	hypoxia or hypercapnia	
J9601	Acute respiratory failure with	Respiratory
	hypoxia	
J9602	Acute respiratory failure with	Respiratory
	hypercapnia	
J9690	Respiratory failure, unsp, unsp w	Respiratory
	hypoxia or hypercapnia	
J9691	Respiratory failure, unspecified	Respiratory
	with hypoxia	
J9692	Respiratory failure, unspecified	Respiratory
Dacaa	with hypercapnia	Dessionles
R0600	Dyspnea, unspecified	Respiratory
R0609	Other forms of dyspnea	Respiratory
R092	Respiratory arrest	Respiratory
J1289	Other viral pneumonia	Respiratory
R0902	Hypoxemia	Respiratory
J9620	Acute and chr resp failure, unsp	Respiratory
10.024	w hypoxia or hypercapnia	
J9621	Acute and chronic respiratory	Respiratory
10(22	failure with hypoxia	Despireter
J9622	Acute and chronic respiratory failure with hypercaphia	Respiratory
R0603	Acute respiratory distress	Respiratory
R0602	Shortness of breath	Respiratory
N170	Acute kidney failure with tubular	renal_failure
1170	necrosis	renal_ianare
N171	Acute kidney failure with acute	renal_failure
	cortical necrosis	<u>renui_</u> runure
N172	Acute kidney failure with	renal failure
	medullary necrosis	
N178	Other acute kidney failure	renal failure
N179	Acute kidney failure, unspecified	 renal failure
K7111	Toxic liver disease with hepatic	hepatic failure
	necrosis, with coma	
K7200	Acute and subacute hepatic	hepatic_failure
	failure without coma	
K7201	Acute and subacute hepatic	hepatic_failure
	failure with coma	
К7290	Hepatic failure, unspecified	hepatic_failure

Severe COVII	D-19	
ICD-10-CM	Description	Туре
	without coma	
K7291	Hepatic failure, unspecified with	hepatic_failure
	coma	
K762	Central hemorrhagic necrosis of	hepatic_failure
	liver	
K763	Infarction of liver	hepatic_failure
D65	Disseminated intravascular	coagulation
	coagulation	
D688	Other specified coagulation	coagulation
	defects	
D689	Coagulation defect, unspecified	coagulation
D6951	Posttransfusion purpura	coagulation
DCOEO	Otherserver	
D6959	Other secondary	coagulation
D696	thrombocytopenia	congulation
F05	Thrombocytopenia, unspecified Delirium due to known	coagulation
FUD	physiological condition	cns_failure
G931	Anoxic brain damage, not	cns failure
0551	elsewhere classified	clis_lallure
G9340	Encephalopathy, unspecified	cns_failure
G9341	Metabolic encephalopathy	cns_failure
G9349	Other encephalopathy	cns_failure
R4020	Unspecified coma	cns failure
1462	Cardiac arrest due to underlying	cardiovascular failure
	cardiac condition	-
1468	Cardiac arrest due to other	cardiovascular_failure
	underlying condition	
1469	Cardiac arrest, cause unspecified	cardiovascular_failure
1951	Orthostatic hypotension	cardiovascular_failure
19589	Other hypotension	cardiovascular_failure
1959	Hypotension, unspecified	cardiovascular_failure
R031	Nonspecific low blood-pressure	cardiovascular_failure
	reading	
R570	Cardiogenic shock	cardiovascular_failure
R571	Hypovolemic shock	cardiovascular_failure
R578	Other shock	cardiovascular_failure
R579	Shock, unspecified	cardiovascular_failure
R6520	Severe sepsis without septic	severe_sepsis
	shock	

Severe COVID-19		
ICD-10-CM	Description	Туре
R6521	Severe sepsis with septic shock	septic_shock
T8112XA	Postprocedural septic shock, initial encounter	septic_shock

Demographic Variables

Dataset Segment:	Demographic Variables
Data Element Name:	Admission Datetime
Template Variable:	admission_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Indicates the date and time that the patient was admitted to inpatient status at the hospital.

Codes and Values:

- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm is NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00
- Cannot have been after *Discharge Datetime*.
- Observation only cases and ED only cases that do not progress to an inpatient admission may use the *Arrival Datetime* as admission date and time.
- If there is a difference between arrival to inpatient floor and the written admission order, report the time the admission order was written.

Demographic Variables

Data Element Name:	Arrival Datetime
Template Variable:	<mark>arrival_dt</mark>
Format – Length:	Datetime – 16
Mandatory:	<mark>Yes</mark>

Description:

Indicates the earliest documented date and time the patient arrived at the hospital.

Codes and Values:

- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm is NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00
- Report earliest date and time the patient arrived at the ED, at the nursing floor, for observation, or as a direct admit to the cath lab.
- The arrival date and time may differ from the Admission Datetime.
- Cannot be after the *Discharge Datetime*.
- Observation Status:
 - If the patient was admitted to observation from an outpatient setting of the hospital, use the date and time the patient arrives at the ED or on the floor of observation care as the arrival date and time.
 - If the patient was admitted to observation from the ED of the hospital, use the date and time the patient arrived at the ED as the *Arrival Datetime*.
- <u>Direct Admits:</u>
 - If the patient is a "Direct Admit" to the cath lab, use the earliest date and time the patient arrived at the cath lab (or cath lab staging/holding area) as the *Arrival Datetime*.
 - If the patient is a "Direct Admit" to acute inpatient or observation, use the earliest date and time the patient arrived at the nursing floor or in observation as the *Arrival Datetime*.

- If the patient was transferred from your hospital's satellite/free-standing ED or from another hospital within your hospital's system (as an inpatient or ED patient) and there is one medical record for the care provided at both facilities, use the *Arrival Datetime* at the first facility.
- The *Arrival Datetime* can be obtained from the time period that the patient was an ED patient.

Demographic Variables

Data Element Name:	Date of Birth
Template Variable:	date_of_birth
Format – Length:	Date — 10
Mandatory:	Yes

Description:

Indicates the date of birth of the patient.

Codes and Values:

- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 - 3. Example: November 3, 1959 = 1959-11-03
- Date of Birth cannot be after Admission Datetime.
- Patient age at admission should be used to determine reporting to the adult or the pediatric database. If a patient is observation only or ED only, please use patient age at the time of arrival for determination of the adult or pediatric database inclusion.
- Data for all patients who are 21 years of age or older are to be reported into the adult NYSDOH database.
 - Patients under 21 as of their admission date (arrival date if not admitted) will be rejected and required for submission to the pediatric sepsis data file.

Demographic Variables

Data Element Name:Discharge DatetimeTemplate Variable:discharge_dtFormat – Length:Datetime — 16Mandatory:Yes

Description:

Indicates the date and time that the patient was discharged from the hospital, left against medical advice, or expired.

Codes and Values:

- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00
- Cannot precede 2014-04-01 00:00.
- Cannot precede Admission Datetime or Arrival Datetime.
- If the time of death and administrative discharge date and times are not the same, use the time of death for *Discharge Datetime*.
- For a patient who is discharged from one unit/department to another unit/department within the same facility, the final discharge from the facility is what should be reported for *Discharge Datetime*. Do not use discharges from internal transfers, since these are not actually separate hospital admissions – the entire period should be submitted as one record. This applies even when the internal transfers are billed separately.

Dataset Segment:	Demographic Variables
Data Element Name:	Discharge Status
Template Variable:	discharge_status
Format – Length:	Enumerated – 2
Mandatory:	<mark>Yes</mark>

Indicates the code that best represents the patient's destination after discharge from the hospital.

Codes and Values:

- 01 = Discharge to Home or Self Care (Routine Discharge). Includes discharge to home; home on oxygen if DME only; any other DME only; group home, foster care, independent living and other residential care arrangements; outpatient programs, such as partial hospitalization or outpatient chemical dependency programs.
- 02 = Discharged/transferred to a Short-Term General Hospital for Inpatient Care.
- 03 = Discharged/transferred to Skilled Nursing Facility (SNF) with Medicare Certification in anticipation of Skilled Care. Medicare indicates that the patient is discharged/transferred to a Medicare certified nursing facility. For hospitals with an approved swing bed arrangement, use Code 61 Swing Bed. For reporting other discharges/transfers to nursing facilities see 04 and 64.
- 04 = Discharged/transferred to a Facility that Provides Custodial or Supportive Care. This is used to designate patients that are discharged/transferred to a nursing facility with neither Medicare nor Medicaid certification and for discharges/transfers to Assisted Living Facilities.
- 05 = Discharged/transferred to a Designated Cancer Center or Children's Hospital.
- 06 = Discharged/transferred to Home under Care of Organized Home Health Service Organization in Anticipation of Covered Skilled Care. Report this code when the patient is discharged/transferred to home with a written plan of care (tailored to the patient's medical needs) for home care services. Not used for home health services provided by a DME supplier or from a Home IV provider for home IV services.
- 07 = Left against medical advice or discontinued care.
- 09 = Admitted as an Inpatient to this Hospital. Patient admitted to the same short-term medical or specialty hospital where the hospital-based ambulatory surgery service was performed (excluding chronic disease hospitals).
- 20 = Expired.
- 21 = Discharged/transferred to Court/Law Enforcement.
- 50 = Hospice Home.
- 51 = Hospice Medical Facility (Certified) Providing Hospice Level of Care.
- 61 = Discharged/transferred to Hospital-Based Medicare Approved Swing Bed.
- 62 = Discharged/transferred to an Inpatient Rehabilitation Facility (IRF), including Rehabilitation Distinct Part Unit of a hospital.

- 63 = Discharged/transferred to a Medicare Certified Long Term Care Hospital (LTCH).
- 64 = Discharged/transferred to a Nursing Facility Certified under Medicaid but not certified under Medicare.
- 65 = Discharged/transferred to a Psychiatric Hospital or Psychiatric Distinct Part Unit of a Hospital.
- 66 = Discharged/transferred to a Critical Access Hospital (CAH).
- 69 = Discharged/transferred to a Designated Disaster Alternative Care Site.
- 70 = Discharged/transferred to another Type of Health Care Institution not defined Elsewhere in this Code List.
- 81 = Discharged to Home or Self Care with a Planned Acute Care Hospital Inpatient Readmission.
- 82 = Discharged/transferred to a Short-Term General Hospital for Inpatient Care with a Planned Acute Care Hospital Inpatient Readmission.
- 83 = Discharged/transferred to Skilled Nursing Facility (SNF) with Medicare Certification with a Planned Acute Care Hospital Inpatient Readmission.
- 84 = Discharged/transferred to a Facility that Provides Custodial or Supportive Care with a Planned Acute Care Hospital Inpatient Readmission.
- 85 = Discharged/transferred to a Designated Cancer Center or Children's Hospital with a Planned Acute Care Hospital Inpatient Readmission.
- 86 = Discharged/transferred to Home under Care of Organized Home Health Service Organization with a Planned Acute Care Hospital Inpatient Readmission.
- 87 = Discharged/transferred to Court/Law Enforcement with a Planned Acute Care Hospital Inpatient Readmission.
- 88 = Discharged/transferred to a Federal Health Care Facility with a Planned Acute Care Hospital Inpatient Readmission.
- 89 = Discharged/transferred to Hospital-Based Medicare Approved Swing Bed with a Planned Acute Care Hospital Inpatient Readmission.
- 90 = Discharged/transferred to an Inpatient Rehabilitation Facility (IRF) including Rehabilitation Distinct Part Units of a Hospital with a Planned Acute Care Hospital Inpatient Readmission.
- 91 = Discharged/transferred to a Medicare Certified Long Term Care Hospital (LTCH) with a Planned Acute Care Hospital Inpatient Readmission.
- 92 = Discharged/transferred to a Nursing Facility Certified under Medicaid but not Certified under Medicare with a Planned Acute Care Hospital Inpatient Readmission.
- 93 = Discharged/transferred to a Psychiatric Hospital or Psychiatric Distinct Part Unit of a Hospital with a Planned Acute Care Hospital Inpatient Readmission.
- 94 = Discharged/transferred to a Critical Access Hospital (CAH) with a Planned Acute Care Hospital Inpatient Readmission.
- 95 = Discharged/transferred to another Type of Health Care Institution not Defined Elsewhere in this Code List with a Planned Acute Care Hospital Inpatient Readmission.

Dataset Segment:	Demographic Variables
Data Element Name:	Ethnicity
Template Variable:	ethnicity
Format – Length:	<mark>Set – maximum 5 codes</mark>
Mandatory:	Yes

Indicates the code that best describes the ethnicity of the patient from the electronic health record (EHR).

Codes and Values:

Examples: E1 = SPANISH/HISPANIC ORIGIN E1.04.004 = Colombian E2 = NOT HISPANIC OR LATINO E9 = UNKNOWN

- If reporting multiple ethnicity codes (up to 5 codes), separate each code using a colon (e.g. "E1.02: E1.04" is Mexican and South American).
- Multiple ethnicity codes within the same heading are expected as there might be many different origins within a heading (e.g. "E1.02.001 Mexican American" and "E1.02.002 Mexicano" are within the same heading "E1.02 Mexican"). However we would not expect a selection of codes within any two headings of "E1 SPANISH/HISPANIC ORIGIN", "E2 NOT HISPANIC OR LATINO", and "E9 UNKNOWN".
- To obtain the full list of codes, please refer to the following link to the SPARCS code set:
- SPARCS(RR-Race and Ethnicity Codes, Source: Race and Ethnicity Code Set -Version 1.0): https://www.health.ny.gov/statistics/sparcs/sysdoc/apprr.htm

Demographic Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Facility Identifier facility_identifier Varchar – 6 Yes

Description:

This number is the facility's four to six-digit Permanent Facility Identifier (PFI) assigned by the Department of Health.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

- Must be a valid number as maintained by the NYSDOH.
- Can only contain numbers 0-9.

Data Element Name:GenderTemplate Variable:genderFormat – Length:Enumerated – 1Mandatory:Yes

Description:

Indicates the gender of the patient.

Codes and Values:

M = Male

F = Female

U = Unknown

Notes for Abstraction:

Demographic Variables

Dataset Segment:	Demographic Variables
Data Element Name:	ICD-10-CM Code (n)
Template Variable:	icd_10_cm_code_(n)
Format – Length:	Set — 8
Mandatory:	Yes

All diagnosis codes (primary and secondary) from the final hospital billed codes. There can be up to 25 codes, and each code will have its own variable and POA indicator. The first ICD-10-CM (Code 1) will be the **principal** diagnosis.

Codes and Values:

- These should be reported as 25 individual variables. The variable fields for these will be as follows:
 - The first Data Element will be ICD-10-CM Code 1 with a template variable of icd_10_cm_code1. ICD-10-CM Code 1 is the PRINCIPAL Diagnosis. All other codes will be secondary diagnosis codes;
 - The twentieth Data Element will be ICD-10-CM Code 20 with a template variable of icd_10_cm_code_20.
- Please provide the final hospital billed codes in this field.
- Hospitals may report up to 25 codes and their indicators, including the principal and secondary codes.
- The ICD-10-CM codes would be submitted WITH the appropriate decimal place (AFTER the 3rd character) for each ICD-10-CM code.

Dataset Segment:	Demographic Variables
Data Element Name:	ICD-10-CM POA Indicator (n)
Template Variable:	icd_10_cm_poa_indicator_(n)
Format – Length:	Enumerated — 1
Mandatory:	Yes

Present on Admission (POA) indicator for each ICD-10-CM diagnosis code, aligning with the data element *ICD-10-CM Code (n)*. The first ICD-10-CM POA (Indicator 1) will be the **principal** diagnosis POA indicator.

Codes and Values:

- Y = Present on admission
- N = Not present on admission
- U = No information in the record
- W = Clinically undetermined
- E = Exempt from POA reporting

- These should be reported as 25 individual variables. The variable fields for these will be as follows:
 - The first Data Element will be ICD-10-CM_POA Indicator 1 with a template variable of icd_10_cm_poa_indicator_1. ICD-10-CM POA Code 1 is the PRINCIPAL Diagnosis POA indicator. All other codes will be secondary diagnosis POA indicators;
 - The twentieth Data Element will be ICD-10-CM POA Indicator 20 with a template variable of icd_10_cm_ poa_indicator_20.
- Please provide the final hospital billed codes in this field.
- Hospitals may report up to 25 POA indicators.
- Please provide the final hospital billed code's POA indicator in this field. Please ensure it aligns with *ICD-10-CM Code (n)*.
- Hospitals are required to report a POA indicator for each ICD-10-CM Code reported.
 - For example, if there are five (5) ICD-10_CM codes reported then five (5) ICD-10-CM POA indicators will be required in the data submission.

Demographic Variables

Data Element Name:	Insurance Number
Template Variable:	insurance_number
Format – Length:	Varchar – 19
Mandatory:	<mark>Yes</mark>

Description:

Indicates the primary insurance policy identification number for the patient.

Codes and Values:

Notes for Abstraction:

- Allows blanks only if Element Payer is not:
 - Medicare ("C")
 - Medicaid ("D")
 - Insurance Company ("F")
 - Blue Cross ("G")
- Must be alphanumeric (0-9) (a-z, A-Z).
- Special characters are invalid entries.

Facilities are directed to enter the following values:

Payer	Type of Number
Blue Cross	Enter the information depending on specific Blue Cross Plan needs and contract requirement.
CHAMPUS	Enter the information depending on CHAMPUS regulations.
Medicaid	Enter Medicaid Client Identification Number (CIN) of the insured or case head
	Medicaid number shown on the Medicaid Identification Card.
Medicare	Enter the patient's Medicare HIC number as shown on the Health Insurance Card,
	Certificate of Award, Utilization Notice, Temporary Eligibility Notice, and Hospital
	Transfer Form or as reported by the Social Security Office.

For all other payer types (commercial insurers, etc.) enter the insured's unique number assigned by the payer.

Dataset Segment:	Demographic Variables
Data Element Name:	Medical Record Number
Template Variable:	medical_record_number
Format – Length:	Varchar – 17
Mandatory:	Yes

Indicates the number used by the hospital's Medical Records Department to identify the patient's permanent medical record file. This number is not the same as the Patient Control Number.

Codes and Values:

- Must not equal zero or blanks.
- Must be alphanumeric (0-9) (a-z, A-Z).
- Special characters are invalid entries.

Dataset Segme	nt:	Demographic Variables	
<mark>Data Element N</mark>	ame:	<mark>Other Payer</mark>	
<mark>Template Varia</mark>	<mark>ole:</mark>	<mark>other_payer</mark>	
<mark>Format – Lengt</mark> l	<mark>ı:</mark>	<mark>Varchar — 50</mark>	
Mandatory:		No	

Indicate the other payers for this hospitalization. This aligns with Payer source E and/or I.

Codes and Values:

- If either E or I is reported under Payer, then Other Payer must be completed.
- If multiple other payers are to be reported, each payer will be separated by a colon (:).
- Include a code and a description if a code is captured in your EHR.

Dataset Segment:	Demographic Variables
Data Element Name:	Patient Control Number
Template Variable:	patient_control_number
Format – Length:	Varchar – 20
Mandatory:	Yes

Indicates the patient's unique number assigned by the provider to facilitate retrieval of individual financial and clinical records and posting of payment.

Codes and Values:

- Must not equal zero or blanks.
- Must be alphanumeric (0-9) (a-z, A-Z).
- Special characters are invalid entries.

Demographic Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Patient Zip Code of Residence patient_zip_code_of_residence Varchar – 10 Yes

Description:

Indicates the patient's 9-digit zip code of residence.

Codes and Values:

- Format should be xxxxx-xxxx
- If a hospital does not have the four-digit extension to the zip code, then the five-digit zip code should be reported followed by 0000 in the extension (e.g., 11201-0000).
- Should only consist of numbers 0-9.

Dataset Segment:	Demographic Variables
Data Element Name:	Payer
Template Variable:	payer
Format – Length:	<mark>Set – maximum 3 codes</mark>
Mandatory:	Yes

Indicate the codes that identify the payers for this hospitalization. Provide the primary payer first.

Codes and Values:

- A = Self-Pay
- B = Workers' Compensation
- C = Medicare
- D = Medicaid
- E = Other Federal Program
- F = Insurance Company
- G = Blue Cross
- H = CHAMPUS
- I = Other Non-Federal Program
- J = Disability
- K = Title V
- L = Unknown

- Report up to 3 payers.
- If either E or I is reported, then Other Payer must be completed.
- Each payer will be separated by a colon (:).
- The <u>PRIMARY</u> payer must be listed first.
 - o Example:
 - Workers' Compensation as primary payer and Disability: B:J
 - Blue Cross as primary payer, Insurance Company, Other Federal Program: G:F:E

Dataset Segment:	Demographic Variables
Data Element Name:	Race
Template Variable:	race
Format – Length:	<mark>Set – maximum 56 codes</mark>
Mandatory:	Yes

Indicates the code that best describes the race of the patient based on the electronic health record.

Codes and Values:

Examples: R2 = Asian R2.01 = Asian Indian R5 = White

- If reporting multiple race codes, separate each code using a colon (e.g. "R2.12: R2.01" is Korean and Asian Indian).
- To obtain the full list of codes, please refer to the following link to the SPARCS code set:
- SPARCS(RR-Race and Ethnicity Codes, Source: Race and Ethnicity Code Set -Version 1.0): https://www.health.ny.gov/statistics/sparcs/sysdoc/apprr.htm

Demographic Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Source of Admission source_of_admission Enumerated – 1 Yes

Description:

Indicates the code that best describes the patient's origin before coming to the hospital.

Codes and Values:

- 1 = <u>Non-Health Facility Point of Origin</u>: The patient was admitted to this facility from home or from an assisted living facility.
- 2 = <u>Clinic</u>: The patient was referred to this facility as a transfer from a freestanding or nonfreestanding clinic.
- 4 = <u>Transfer from a Hospital (Different Facility</u>): The patient was admitted to this facility as a hospital transfer from an acute care facility where he or she was an inpatient or outpatient.
- 5 = <u>Transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)</u>: The patient was admitted to this facility as a transfer from a SNF or ICF where he/she was a resident.
- 6 = <u>Transfer from Another Health Care Facility</u>: The patient was admitted to this facility as a transfer from another type of health care facility that is not defined elsewhere in this code list.
- 8 = <u>Court/Law Enforcement</u>: The patient was admitted to this facility upon the direction of a court of law or upon the request of a law enforcement agency representative.
- 9 = <u>Information Not Available</u>: The means by which the patient was admitted to this hospital was not known.
- E = <u>Transfer from Ambulatory Surgery Center</u>: The patient was admitted to this facility as a transfer from an ambulatory surgery center.
- F = <u>Transfer from Hospice and is Under a Hospice Plan of Care or Enrolled in a Hospice</u> <u>Program</u>: The patient was admitted to this facility as a transfer from a hospice.

- If a patient is moved from one area of the hospital to another (e.g., from the Emergency Department to the ICU), the patient is not considered a transfer. The patient is considered a transfer when the patient is moved between different hospitals with discharge and admission at each location and separate billing from each location.
- Assisted Living is reported as 1, Non-Health Facility Point of Origin.

Dataset Segment:	Demographic Variables
Data Element Name:	Transferred In
Template Variable:	transferred_in
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates if the patient was received as a transfer from another acute care hospital.

Codes and Values:

<mark>0 = No</mark> 1 = Yes

- Report "1", if a patient was transferred in (i.e., received from another acute care hospital).
- Report "0", if a patient was not transferred in.

<mark>Datas</mark>	et Segment:	Demographic Variables
	lement Nam <mark>e:</mark>	Transferred Out
<mark>Templ</mark>	<mark>ate Variable:</mark>	transferred_out
<mark>Forma</mark>	<mark>t – Length:</mark>	<mark>Enumerated – 1</mark>
<mark>Mand</mark>	atory:	<mark>Yes</mark>

Indicates if the patient was transferred out to another acute care hospital.

Codes and Values:

<mark>0 = No</mark> 1 = Yes

- Report "1", if a patient was transferred out (i.e., transferred/discharged to another acute care hospital).
- Report "0", if a patient was not transferred out.

Demographic Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Transfer Facility Identifier Receiving transfer_facility_id_receiving Varchar – 6 No

Description:

If your hospital received a transfer patient from an acute care hospital, report the hospital PFI from which you received that patient. This is the transferring facility's four to six-digit Permanent Facility Identifier (PFI) assigned by the Department of Health.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Notes for Abstraction:

- Must be a valid number as maintained by the NYSDOH.
- Must only contain numbers 0-9.
- When receiving a patient from an out-of-state facility, please submit the two-digit state identifier https://www2.census.gov/geo/docs/reference/state.txt to represent the transfer facility state. This is ONLY to be used when patients are received from an out of state hospital, therefore the code for New York will not be accepted for data submission. For example, a patient received from a Connecticut hospital is submitted with the *transfer_facility_id_receiving* of 09.

To find a hospital PFI, please visit:

https://www.health.ny.gov/statistics/sparcs/reports/compliance/pfi facilities.htm

Demographic Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Transfer Facility Identifier Sending transfer_facility_id_sending Varchar – 6 No

Description:

If your hospital is transferring a patient to another acute care hospital, report the hospital's PFI to which you are sending the patient. This number is the transfer sending facility's four to six-digit Permanent Facility Identifier (PFI) assigned by the Department of Health.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Notes for Abstraction:

- Must be a valid number as maintained by the NYSDOH.
- Must only contain numbers 0-9.
- When transferring a patient to an out-of-state facility, please submit the two-digit state identifier https://www2.census.gov/geo/docs/reference/state.txt to represent the transfer facility state. This is ONLY to be used when patients are transferred out of state therefore the code for New York will not be accepted for data submission. For example, a patient transferred to a Connecticut hospital is submitted with the *transfer_facility_id_sending* of 09.

To find a hospital PFI, please visit:

https://www.health.ny.gov/statistics/sparcs/reports/compliance/pfi facilities.htm

Data Element Name: Template Variable: Format – Length: Mandatory:

Demographic Variables

Transfer Facility Name Receiving transfer_facility_nm_receiving Varchar – 50 No

Description:

If your hospital received a patient as a transfer from another acute care hospital, report the hospital name from which you received that patient.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Notes for Abstraction:

Report when Transfer Facility Identifier Receiving is not available.

Data Element Name: Template Variable: Format – Length: Mandatory:

Demographic Variables

Transfer Facility Name Sending transfer_facility_nm_sending Varchar – 50 No

Description:

<u>If your hospital is transferring a patient to an acute care hospital, report the hospital's name to</u> which you are sending the patient.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Notes for Abstraction:

• Report when Transfer Facility Identifier Sending is not available.

Demographic Variables

Data Element Name:Unique Personal IdentifierTemplate Variable:unique_personal_identifierFormat – Length:Varchar – 10Mandatory:Yes

Description:

A composite field comprised of portions of the patient last name, first name, and social security number.

Codes and Values:

Included below are the individual components of this data element.

- 1. **"First 2" and "Last 2" characters of the Patient's Last Name**. The birth name of the patient is preferable if it is available on the facility's information system.
- 2. "First 2" characters of the Patient's First Name.
- 3. "Last 4" digits of the Patient's Social Security Number.

NOTE: This data element is not to be confused with *Patient Control Number*, which provides linkage of all record types containing patient-related data for a specific discharge.

Notes for Abstraction:

First and Last Name Components: Must be <u>UPPERCASE</u> alpha characters (A-Z). If the last name is less than 4 characters, the first two and last two characters are used even if some characters are repeated.

Included below are examples of how to report some unusual scenarios: A three-character last name, a two-character last name, a name with junior, a one character first name, a last name with an apostrophe, and a hyphenated last name.

- Joe Tan would be reported as TAANJO
- Bill Su Jr. would be reported as SUSUBI
- E John Smith would be reported as SMTHEE
- Bob O'Brien would be reported as OBENBO
- Sue Jones-Davis would be reported as JOISSU

Social Security Number Component: Must be numeric. If no Social Security Number is available, this sub-field must be zeroes (e.g. TAANJO0000).

Joe Tan with Social Security Number 123-456-7890 would be reported as TAANJO7890

Comorbidity/Risk Factor (POA) Variables

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Acute Cardiovascular Conditions
Template Variable:	acute_cardiovascular_conditions
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient had an <u>acute</u> cardiovascular event present at admission/arrival. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

1= Myocardial Infarction2= Ischemic Stroke/Hemorrhagic Stroke/Transient Ischemic Attack (TIA)0= No Acute Cardiovascular Condition

- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2
- Please see Appendix 1A for a list of applicable ICD-10-CM codes.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: AIDS/HIV Disease aids_hiv_disease Enumerated – 1 <mark>Yes</mark>

Description:

Indicates that the patient has AIDS or an HIV infection.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1B for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Altered Mental Status altered_mental_status Enumerated – 1 <mark>Yes</mark>

Description:

Indicates that the patient has an altered mental status.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1C for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Asthma
Template Variable:	asthma
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient has asthma.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1D for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Data Element Name: Template Variable: Format – Length: Mandatory:

Description:

Indicates that the patient has chronic liver disease.

Codes and Values:

0 = No 1 = Yes

Notes for Abstraction:

- Please see Appendix 1E for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Chronic Liver Disease chronic_liver_disease Enumerated – 1 Yes

Data Element Name: Template Variable: Format – Length: Mandatory:

Description:

Indicates that the patient has chronic renal failure.

Codes and Values:

0 = No 1 = Yes

Notes for Abstraction:

- Please see Appendix 1F for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Chronic Renal Failure chronic_renal_failure Enumerated – 1 <mark>Yes</mark>

Comorbidity/Risk Factor (POA) Variables

Chronic Respiratory Failure chronic_respiratory_failure Enumerated – 1 Yes

Data Element Name: Template Variable: Format – Length: Mandatory:

Description:

Indicates that the patient has chronic respiratory failure.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1G for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	et Segment:	Segn	et	Datas
------------------	-------------	------	----	-------

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: Mandatory:

Description:

Indicates that the patient has coagulopathy.

Codes and Values:

0 = No 1 = Yes

Notes for Abstraction:

- Please see Appendix 1H for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Coagulopathy coagulopathy Enumerated – 1 <mark>Yes</mark>

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark>

Comorbidity/Risk Factor (POA) Variables

Congestive Heart Failure congestive_heart_failure Enumerated – 1 <mark>Yes</mark>

Description:

Indicates that the patient has congestive heart failure.

Codes and Values:

0 = No 1 = Yes

I = Yes

- Please see Appendix 1I for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	COPD
Template Variable:	copd
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient has chronic obstructive pulmonary disease (COPD).

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1J for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Dementia
Template Variable:	dementia
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient has dementia.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1K for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Diabetes
Template Variable:	diabetes
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient has diabetes.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1L for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Dialysis Comorbidity
Template Variable:	dialysis_comorbidity
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient arrived at the hospital already receiving dialysis. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

- 0 = No
- 1 = Yes

- Please see Appendix 1M for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> History of COVID -19 history_of_covid Enumerated – 1 Yes

Description:

Indicates that the patient has a history of a positive COVID-19 test.

Codes and Values:

0 = No 1 = Yes

- Report "1" when the patient has a history of a positive COVID-19 test prior to arrival at the hospital. There is not a time limit on reporting the test as positive. If there is a positive test, even if later followed by a negative test, then report the positive test date.
- SARS-Cov-2 LOINC codes can be downloaded to a csv. This file can be found here: <u>https://loinc.org/sars-cov-2-and-covid-19/</u>
- These codes are not static and are updated regularly; therefore, hospitals should take care to use the most current list of codes to capture COVID-19 testing.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> History of COVID-19 Datetime history_of_covid_dt Datetime – 16 No

Description:

Indicates the date and time of the positive *History of COVID-19* test.

Codes and Values:

- If there is more than one positive COVID-19 test, report the earliest positive test.
- <u>Formatting:</u>
 - Format must be YYYY-MM-DD hh:mm

 YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> History of Other Cardiovascular Disease history_of_other_cvd Set – maximum 4 codes Yes

Description:

Indicates the patient's history of other cardiovascular disease.

Codes and Values:

- 1 = Coronary heart disease (e.g. angina pectoris, coronary atherosclerosis)
- 2 = Peripheral artery disease
- 3 = Valve disorder
- 4 = Cerebrovascular disease
- 0 = No history of coronary heart disease, peripheral artery disease, valve disorder or cerebrovascular disease

- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2:3
- History of (not acute presentation)
- Please see Appendix 1N for a list of applicable ICD-10-CM codes.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Hypertension
Template Variable:	hypertension
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient has hypertension.

Codes and Values:

0 = No

1 = Yes

- Please see Appendix 10 for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Immunocompromising immunocompromising Enumerated – 1 Yes

Description:

Indicates that the patient has an immunocompromising disease/illness.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1P for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Lymphoma Leukemia Multiple Myeloma
Template Variable:	lymphoma_leukemia_multi_myeloma
Format – Length:	Enumerated – 1
Mandatory:	<mark>Yes</mark>

Indicates that the patient has malignant neoplasm of lymphatic and hematopoietic tissue including those neoplasms which may be in clinical remission.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1Q for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables	
Data Element Name:	Mechanical Ventilation Comorbidity	
Template Variable:	mechanical_vent_comorbidity	
Format – Length: <mark>Mandatory:</mark>	Enumerated – 1 <mark>Yes</mark>	

Indicates that the patient arrived at the hospital on mechanical ventilation. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No

1 = Yes

- Please see Appendix 1R for applicable ICD-10-CM code.
- Report "1", if the patient has the ICD-10-CM code listed in the referenced appendix.
- Report "0", if the patient does not have the ICD-10-CM code listed in the referenced appendix.

Data Element Name:

Template Variable:

Format – Length: Mandatory:

Comorbidity/Risk Factor (POA) Variables

Medication Anticoagulation medication_anticoagulation Enumerated – 1 Yes

Description:

Indicates that the patient takes anticoagulation medications at home/prior to admission.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1S for a list of applicable medications and NDC codes.
- Report "1", if the patient has one or more of the medications or NDC codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the medications or NDC codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Medication Immune Modifying
Template Variable:	medication_immune_modifying
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient is taking disease modifying medications and therapies (drugs and biologics) for collagen diseases, corticosteroids, chemotherapeutic agents through any modality (oral, IV, IM, etc.) known to specifically adversely impact the function of the immune system as the primary therapeutic goal or as an unintended side effect, including steroids (excluding inhaled or topical steroids) and chemotherapy at time of admission.

Codes and Values:

_ ...

- Please see Appendix 1T for a list of applicable medications and NDC codes.
- Report "1", if the patient has one or more of the medications or NDC codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the medications or NDC codes listed in the referenced appendix.

Data Element Name: Template Variable: Format - Length: Mandatory:

Description:

Indicates that the patient has metastatic cancer.

Codes and Values:

0 = No 1 = Yes

Notes for Abstraction:

- Please see Appendix 1U for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Metastatic Cancer metastatic_cancer Enumerated – 1 Yes

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Obesity
Template Variable:	obesity
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient is obese (measured as a body mass index (BMI) of 30 or higher).

Codes and Values:

0 = No 1 = Yes

- Combination of ICD-10-CM and/or BMI values from the electronic health record (EHR). Please use the first value upon admission/arrival or the earliest value.
- Please see Appendix 1V for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "1", if the patient has a BMI value of 30 or higher in the EHR even if they do not have one of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix or a BMI value of 30 or higher in the EHR.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Patient Care Considerations patient_care_considerations Set – maximum 3 codes Yes

Description:

Indicates whether the patient has a Do Not Resuscitate (DNR), Do Not Intubate (DNI) or comfort care order at any time during the hospital encounter.

Codes and Values:

1 = DNR 2 = DNI 3 = Comfort Care 0 = None

- Report all that apply.
- Each payer will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2:3

Dataset Segment:	Comorbidity/Risk Factor (POA) Variables
Data Element Name:	Patient Care Considerations Date
Template Variable:	patient_care_considerations_date
Format – Length:	Date – 10
Mandatory:	No

Indicate the earliest date associated with *patient_care_considerations* (i.e., DNR, DNI, and/or Comfort Care).

Codes and Values:

- Format must be YYYY-MM-DD
 - a. YYYY = four-digit year
 - b. MM = two-digit month (01 = January, etc.)
 - c. DD = two-digit day of month (01 through 31)
- Example: November 3, 1959 = 1959-11-03

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Pregnancy Comorbidity pregnancy_comorbidity Enumerated – 1 Yes

Description:

Indicates that the patient has a pregnancy-related comorbidity.

Codes and Values:

0 = No 1 = Yes

- Must be "0" if Pregnancy Status During Hospitalization is "0"
- Please see Appendix 1W for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Comorbidity/Risk Factor (POA) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Pregnancy Status During Hospitalization pregnancy_status Enumerated – 1 Yes

Description:

Indicates the patient is pregnant, in childbirth, or postpartum on arrival to the hospital or during hospitalization.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1X for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "1" if detection of human chorionic gonadotropic (hCG) in the urine or blood test (lab value).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix or does not have detection of human chorionic gonadotropic (hCG) in the urine or blood test (lab value).

Comorbidity/Risk Factor (POA) Variables

Data Element Name:		
Template Variable:		
Format – Length:		
<mark>Mandatory:</mark>		

Smoking Vaping smoking_vaping Enumerated – 1 Yes

Description:

Indicates that the patient is a current smoker and/or a current vaper.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1Y for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Comorbidity/Risk Factor (POA)
Data Element Name:	Tracheostomy on Arrival
Template Variable:	tracheostomy_on_arrival
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates that the patient has a tracheostomy upon admission/arrival to the hospital. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1Z for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Clinical Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark>

Clinical Variables

COVID-19 Exposure covid_exposure Enumerated – 1 Yes

Description:

Indicates the patient has exposure to COVID-19.

Codes and Values:

0 = No Positive COVID-19 exposure

1 = Positive COVID-19 exposure

- Please see Appendix 2A for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Clinical Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> COVID-19 Virus covid_virus Enumerated – 1 <mark>Yes</mark>

Description:

Indicates COVID-19 virus is identified or not identified.

Codes and Values:

- 0 = COVID-19, virus not identified
- 1 = COVID-19, virus identified

- This applies to both present on admission/arrival (POA) or acquired during hospitalization.
- Please see Appendix 2B for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Clinical Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Drug Resistant Pathogen drug_resistant_pathogen Enumerated – 1 Yes

Description:

Indicates that the patient has resistance to an antimicrobial drug.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 2C for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Clinical Variables
Data Element Name:	Flu Positive
Template Variable:	flu_positive
Format – Length:	Enumerated – 1
Mandatory:	<mark>Yes</mark>

Indicates that the patient has a positive flu test present on admission/arrival or during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- POA and/or during hospitalization
- Please see Appendix 2D for a list of applicable codes.
- Report "1", if the patient has one or more of the codes listed in the referenced appendix.
- Report "1", if the patient has a positive influenza virus test (lab value).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix or does not have a positive influenza virus test (lab value).

Data Element Name: Template Variable:

Format – Length: <mark>Mandatory:</mark>

Description:

The suspected source of infection.

Codes and Values:

- 1 = septicemia
- 2 = bacteremia
- 3 = fungal infection
- 4 = peritoneal infection
- 5 = heart infection
- 6 = upper respiratory infection
- 7 = lung infection
- 8 = central nervous system infection
- 9 = gastrointestinal infection
- 10 = genitourinary infection
- 11 =soft tissue infection
- 12 = other infection source
- 13 = unknown

Notes for Abstraction:

- Please see Appendix 2E for a list of applicable ICD-10-CM codes.
- If there is not an identified source of infection as specified in Appendix 2E, then report "unknown".
 - Note that "other infection source" is defined in the ICD-10-CM codes provided in the appendix.
- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 2:9:12

Clinical Variables

Suspected Source of Infection suspected_source_of_infection Set – maximum 12 codes Yes **Treatment (during hospitalization) Variables**

Treatment (during hospitalization) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Dialysis Treatment dialysis_treatment Enumerated – 1 <mark>Yes</mark>

Description:

Indicates that the patient has an order for dialysis during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- Please see Appendix 3A for a list of applicable ICD-10-PCS codes.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced appendix.
- Report "1", if the patient has an order for dialysis in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-PCS codes listed in the referenced appendix or does not have an order for dialysis in the EHR.

Dataset Segment:	Treatment (during hospitalization) Variables
Data Element Name:	During Hospital Anticoagulation
Template Variable:	during_hospital_anticoagulation
Format – Length:	Number – 1
Mandatory:	Yes

Indicates that the patient has an order for anticoagulation medication during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1S for a list of applicable medications and NDC codes.
- Report "1", if the patient has one or more of the medications or NDC codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the medications or NDC codes listed in the referenced appendix.

Dataset Segment:	Treatment (during hospitalization) Variables
Data Element Name:	During Hospital Immune Modifying Medication
Template Variable:	during_hospital_immune_mod_med
Format – Length:	Number – 1
Mandatory:	Yes

Indicates that the patient has an order for immune-modifying medication during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 1T for a list of applicable medications and NDC codes.
- Report "1", if the patient has one or more of the medications or NDC codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the medications or NDC codes listed in the referenced appendix.

Treatment (during hospitalization) Variables

Data Element Name:During Hospital RemdesivirTemplate Variable:during_hospital_remdesivirFormat – Length:Number – 1Mandatory:Yes

Description:

Indicates the patient has an order for remdesivir during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Generic: Remdesivir
- Brand name: Veklury and GS-5734
- Please see Appendix 3B for a list of applicable ICD-10-PCS codes.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced appendix.
- Report "1", if the patient has an order for the medication listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-PCS codes or the medication listed in the referenced appendix.

Dataset Segment:	Treatment (during hospitalization) Variables
Data Element Name:	ECMO
Template Variable:	ecmo
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates the patient has an order for extracorporeal membrane oxygenation (ECMO) during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- Please see Appendix 3C for a list of applicable ICD-10-PCS codes.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced appendix.
- Report "1", if the patient has an order ECMO in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix or does not have an order for ECMO in the EHR.

Treatment (during hospitalization) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: High Flow Nasal Cannula high_flow_nasal_cannula Enumerated – 1 <mark>Yes</mark>

Description:

Indicates the patient has an order for high flow nasal cannula at any time during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Report "1" when the patient has an order for high flow nasal cannula at any time during the hospital encounter.
- Report "0", if the patient does not have an order for high flow nasal cannula at any time during the hospital encounter.

Treatment (during hospitalization) Variables

Mechanical Ventilation Treatment mechanical_vent_treatment Enumerated – 1 Yes

Data Element Name: Template Variable: Format – Length: Mandatory:

Description:

Indicates the patient has an order for mechanical ventilation at any time during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- Please see Appendix 3D for a list of applicable ICD-10-PCS codes.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced appendix.
- Report "1", if the patient has an order for mechanical ventilation in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix or does not have an order for mechanical ventilation in the EHR.

Dataset Segment:	Treatment (during hospitalization) Variables
Data Element Name:	Non-Invasive Positive Pressure Ventilation
Template Variable:	non invasive pos pressure vent
Format – Length:	Enumerated – 1
<mark>Mandatory:</mark>	Yes

Indicates the patient has an order for non-invasive-positive pressure ventilation (CPAP, BiPAP) during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- Please see Appendix 3E for a list of applicable ICD-10-PCS codes.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced appendix.
- Report "1", if the patient has an order for non-invasive positive pressure ventilation in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix or does not have an order for non-invasive positive pressure ventilation in the EHR.

Treatment (during hospitalization) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Vasopressor Administration vasopressor_administration Enumerated – 1 Yes

Description:

Indicates the patient has an order for vasopressors during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 3F for a list of applicable medications and NDC codes.
- Report "1", if the patient has one or more of the medications or NDC codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the medications or NDC codes listed in the referenced appendix.

Outcome (at discharge) Variables

Outcome (at discharge) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Dialysis Outcome dialysis_outcome Enumerated – 1 Yes

Description:

Indicates that the patient is discharged on dialysis.

Codes and Values:

0 = No 1 = Yes

- If there was a patient order to have dialysis at discharge as evidenced by dialysis on the discharge date, report "1."
- Please see Appendix 4A for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has the ICD-10-CM code listed in the referenced appendix on the date of discharge.
- Report "0", if the patient does not have the ICD-10-CM code listed in the referenced appendix on the date of discharge.

Data Element Name:

Template Variable:

Outcome (at discharge) Variables

Mechanical Ventilation Outcome mechanical_vent_outcome Enumerated – 1 Yes

Format – Length: <mark>Mandatory:</mark>

Description:

Indicates the patient is discharged on mechanical ventilation.

Codes and Values:

0 = No 1 = Yes

- If there was a patient order to have mechanical ventilation at discharge as evidenced by mechanical ventilation on the discharge date, report "1".
- Please see Appendix 4B for a list of applicable ICD-10-CM codes. Report "1", if the patient has the ICD-10-CM code listed in the referenced appendix on the date of discharge.
- Report "0", if the patient does not have the ICD-10-CM code listed in the referenced appendix on the date of discharge.

Outcome (at discharge) Variables

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Tracheostomy at Discharge tracheostomy_at_discharge Enumerated – 1 Yes

Description:

Indicates that the patient was discharged with a tracheostomy.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 4C for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has the ICD-10-CM code listed in the referenced appendix on the date of discharge.
- Report "0", if the patient does not have the ICD-10-CM codes listed in the referenced appendix on the date of discharge.

Outcome (during hospitalization) Variables

Outcome (in hospital) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Cardiovascular Outcomes cardiovascular_outcomes Enumerated – 1 Yes

Description:

Indicates the patient had one of the following cardiovascular_outcome(s) during the hospitalization.

Codes and Values:

- 0 = None
- 1 = Acute Coronary Syndrome
- 2 = Ischemic Stroke
- 3 = Myocarditis secondary to COVID-19

- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2:3
- Please see Appendix 4D for a list of applicable ICD-10-CM codes.

Data Element Name: Template Variable: Format – Length: Mandatory:

Outcome (in hospital) Variables

ICU During Hospitalization icu_during_hospitalization Enumerated – 1 Yes

Description:

Indicate if the patient was admitted to the Intensive Care Unit (ICU; MICU; SICU; CCU; Neuro-ICU) during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Report "1", if the patient was admitted at any time to the ICU during the hospital admission.
- Report "0", if the patient was not admitted to the ICU during the hospital admission.

Dataset Segment:	Outcome (in hospital) Variables
Data Element Name:	PE/DVT
Template Variable:	pe_dvt
Format – Length:	Enumerated – 1
Mandatory:	Yes

Indicates the patient had a pulmonary embolism (PE) and/or deep venous thrombosis (DVT) during the hospitalization.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 4E for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Outcome (in hospital) Variables

Data Element Name: Template Variable: Format – Length: Mandatory: Tracheostomy in Hospital tracheostomy_in_hospital Enumerated – 1 Yes

Description:

Indicates that the patient had a tracheostomy during the hospitalization prior to the discharge date.

Codes and Values:

0 = No 1 = Yes

- If the patient received a tracheotomy at arrival or during the hospitalization, report "1."
- Please see Appendix 4F for a list of applicable ICD-10-PCS codes.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced appendix.
- Report "0", if the patient does not have one or more of the ICD-10-PCS codes listed in the referenced appendix.

Severity Variables

Dataset Segment:	Severity Variables
Data Element Name:	aPTT 1
Template Variable:	aptt_1
Format – Length:	String – 4
Mandatory:	No

Indicates the first activated partial thromboplastin time (aPTT) level collected after arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	aPTT 2
Template Variable:	aptt_2
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the second aPTT value collected after arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	aPTT 3
Template Variable:	aptt_3
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the third aPTT level collected after arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	aPTT Max
Template Variable:	aptt_max
Format – Length:	String – 4
Mandatory:	No

Indicates the maximum aPTT value collected after arrival to the hospital.

Codes and Values:

Enter the aPTT level.

- **aPTT 1/2/3/Max** and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- Must be reported to one decimal place (example 19.8). For example, 30.7 or 30.0; place hold with 0.
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.
- If the aPTT level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding the aPTT level results:
 - o 30.48 is rounded to 30.5
 - 45.43 is rounded to 45.4
 - o 61.75 is rounded to 61.8
 - o 55.97 is rounded to 56.0
 - NOT CORRECT: 61.75 is truncated to 61.7 (this should be rounded to 61.8)

Dataset Segment:	Severity Variables
Data Element Name:	aPTT Datetime 1
Template Variable:	aptt_dt_1
Format – Length:	Datetime – 16
Mandatory:	No No

Indicates the date and time of the first aPTT level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	aPTT Datetime 2
Template Variable:	aptt_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second aPTT level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	aPTT Datetime 3
Template Variable:	aptt_dt_3
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	No

Description:

Indicates the date and time of the third aPTT level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name: Template Variable:	aPTT Datetime Max <mark>aptt_dt_max</mark>
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	No No

Indicates the date and time of the maximum aPTT level collected after the patient's arrival to the hospital.

Codes and Values:

- aPTT 1/2/3/Max Datetimes are situational because hospitals may not always have values to report.
 - If any aPTT is reported then the datetime for the aPTT should be reported. For example, if aPTT 1 has a value, then aPTT Datetime 1 should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- This element belongs to Organ Dysfunction Hematologic.
- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic First
Template Variable:	diastolic_1
Format – Length:	Number – 3
Mandatory:	Yes

Indicates the patient's first diastolic blood pressure collected after arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic Second
Template Variable:	diastolic _2
Format – Length:	Number – 3
Mandatory:	No

Description:

Indicate the patient's second diastolic blood pressure collected after arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic Third
Template Variable:	diastolic_3
Format – Length:	Number – 3
Mandatory:	No

Description:

Indicate the patient's third diastolic blood pressure collected after arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic Min
Template Variable:	diastolic _min
Format – Length:	Number – 3
Mandatory:	No

Description:

Indicates the patient's minimum diastolic blood pressure collected after arrival to the hospital.

Codes and Values:

- Diastolic First and corresponding Diastolic First Datetime 1 are mandatory. However, Diastolic Second, Diastolic Third, and Diastolic Min and corresponding datetimes are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - If any Diastolic is reported then the datetime for the Diastolic value should be reported. For example, if *Diastolic Second* has a value, then *Diastolic Second Datetime 2* should not be blank.
- <u>Formatting:</u>
- Format must be a number up to 3 digits.
 - 1. Example:
 - a. Diastolic blood pressure 80mm Hg should be reported as 80
 - b. Diastolic blood pressure 112 Hg should be reported as 112

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic First Datetime 1
Template Variable:	diastolic _dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Indicates the date and time of the first diastolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic Second Datetime 2
Template Variable:	diastolic_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second diastolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic Third Datetime 3
Template Variable:	diastolic _dt_3
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	No

Description:

Indicates the date and time of the third diastolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Diastolic Datetime Min
Template Variable: Format – Length:	<mark>diastolic _dt_min</mark> Datetime – 16
Mandatory:	No

Indicates the date and time of the minimum diastolic blood pressure collected after the patient's arrival to the hospital.

Codes and Values:

- Diastolic First Datetime 1 is mandatory. However, Diastolic Second Datetime 2, Diastolic Third Datetime 3, Diastolic Datetime Min are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - If any Diastolic is reported then the datetime for the Diastolic value should be reported. For example, if *Diastolic Second* has a value, then *Diastolic Second Datetime 2* should not be blank.
- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	INR 1
Template Variable:	inr_1
Format – Length:	String – 4
Mandatory:	No

Indicates the first INR value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR 2
Template Variable:	inr_2
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the second INR level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR 3
Template Variable:	inr_3
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the third INR level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR Max
Template Variable:	inr_max
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the maximum INR level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the INR level

- *INR 1/2/3/Max* and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- Must be reported to one decimal place (example 1.2 or 11.5).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.
- If the INR level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding INR level results:
 - o 2.48 is rounded to 2.5
 - o 11.75 is rounded to 11.8
 - 2.97 is rounded to 3.0
 - NOT CORRECT: 11.75 is truncated to 11.7 (this should be rounded to 11.8)

Dataset Segment:	Severity Variables
Data Element Name:	INR Datetime 1
Template Variable:	inr_dt_1
Format – Length:	Datetime – 16
Mandatory:	No

Indicates the date and time of the first INR level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR Datetime 2
Template Variable:	inr_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second INR level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR Datetime 3
Template Variable:	inr_dt_3
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the third INR collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR Datetime Max
Template Variable: Format – Length: <mark>Mandatory:</mark>	<mark>inr_dt_max</mark> Datetime – 16 <mark>No</mark>

Description:

Indicates the date and time of the maximum INR level collected after the patient's arrival to the hospital.

Codes and Values:

- INR 1/2/3/Max Datetimes are situational because hospitals may not always have values to report.
 - If any INR is reported then the datetime for the INR value should be reported. For example, if *INR* 1 has not value, *INR Datetime* 1 should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- This element belongs to Organ Dysfunction Hematologic.
- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level 1
Template Variable:	lactate_level_1
Format – Length:	String – 4
Mandatory:	No

Indicates the first lactate level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level 2
Template Variable:	lactate_level_2
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the second lactate level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level 3
Template Variable:	lactate_level_3
Format – Length:	String – -4
Mandatory:	No.

Description:

Indicates the third lactate level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level Max
Template Variable:	lactate_level_max
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the maximum lactate level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual lactate level using the mmol/L value. Convert from mg/dL if needed.

• Values might range from 0 to 9; numbers higher than nine may indicate the value has not been converted to mmol/L.

- Lactate Level 1/2/3/Max and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- Must be reported to one decimal place (example 19.8).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.
- If the lactate level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding lactate level results:
 - o 7.81 is rounded to 7.8
 - o 7.85 is rounded to 7.9
 - o 7.23 is rounded to 7.2
 - 7.97 is rounded to 8.0
 - NOT CORRECT: 7.85 is truncated to 7.8 (this should be rounded to 7.9)

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level Datetime 1
Template Variable:	lactate_level_dt_1
Format – Length:	Datetime – 16
Mandatory:	No

Indicates the date and time of the first lactate level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level Datetime 2
Template Variable:	lactate_level_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second lactate level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level Datetime 3
Template Variable:	lactate_level_dt_3
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	<mark>No</mark>

Description:

Indicates the date and time of the third lactate level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Lactate Level Datetime Max
Template Variable:	<mark>lactate_level_dt_max</mark>
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	No

Indicates the date and time of the maximum lactate level collected after the patient's arrival to the hospital.

- Lactate Level 1/2/3/Max Datetimes are situational because hospitals may not always have values to report.
 - If any Lactate Level is reported then the datetime for the Lactate Level value should be reported. For example, if *Lactate Level 1* has a value, *Lactate Level Datetime 1* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - 2. YYYY = four-digit year
 - MM = two-digit month (01 = January, etc.) DD = two-digit day of month (01 through 31) hh = two digits of hour (00 through 23) (am/pm NOT allowed) mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:

Data Element Name: Template Variable: Format – Length: Mandatory:

Severity Variables

Organ Dysfunction CNS organ_dysfunc_cns Enumerated – 1 Yes

Description:

Indicates that the patient has central nervous system (CNS) organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 5A for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Hepatic Arrival
Template Variable:	organ_dysfunc_hepatic_arrival
Format – Length:	String – 4
Mandatory:	No

Indicates the first bilirubin level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Hepatic Max
Template Variable:	organ_dysfunc_hepatic_max
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the maximum bilirubin level collected after arrival to the hospital.

Codes and Values:

Enter the actual bilirubin level. Convert the units to mg/dL if needed.

- **Organ Dysfunction Hepatic Arrival/Max** and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- Must be reported to one decimal place (example 2.8).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.
- If the bilirubin level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding bilirubin level results:
 - o 2.51 is rounded to 2.5
 - o .75 is rounded to .8
 - 1.97 is rounded to 2.0
 - **NOT CORRECT:** .75 is truncated to .7 (this should be rounded to .8)

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Hepatic Arrival Datetime
Template Variable:	organ_dysfunc_hepatic_arrival_dt
Format – Length: <mark>Mandatory:</mark>	Datetime – 16 <mark>No</mark>

Indicates the date and time of the first bilirubin collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Hepatic Max Datetime
Template Variable:	organ_dysfunc_hepatic_max_dt
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the maximum bilirubin level collected after arrival to the hospital.

Codes and Values:

- Organ Dysfunction Hepatic Arrival/Max Datetimes are situational because hospitals may not always have values to report.
 - If any Organ Dysfunction Hepatic is reported then the datetime for Organ Dysfunction Hepatic should be reported. For example, if Organ Dysfunction Hepatic Arrival has a value, Organ Dysfunction Hepatic Arrival Datetime should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)

hh = two digits of hour (00 through 23) (am/pm NOT allowed) mm = two digits of minute (00 through 59)

- 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
- 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Renal Arrival

Template Variable: Format – Length: Mandatory:

organ dysfunc renal arrival String – 4 No

Description:

Indicates the first creatinine level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Renal Max
Template Variable:	organ_dysfunc_renal_max
Format – Length:	String – 4
Mandatory:	No

Description:

Indicates the maximum creatinine level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual creatinine level. Convert the units to mg/dL if needed.

- Organ Dysfunction Renal Arrival/Max and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- Must be reported to one decimal place (example 2.8).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.
- If the creatinine level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding creatinine level results:
 - o 2.81 is rounded to 2.8
 - o 1.75 is rounded to 1.8
 - o 1.42 is rounded to 1.4

- o 2.97 is rounded to 3.0
- NOT CORRECT: 1.75 is truncated to 1.7 (this should be rounded to 1.8)

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Renal Arrival Datetime
Template Variable:	organ_dysfunc_renal_arrival_dt
Format – Length: <mark>Mandatory:</mark>	Datetime – 16 <mark>No</mark>

Indicates the date and time of the first creatinine level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Organ Dysfunction Renal Max Datetime
Template Variable:	organ_dysfunc_renal_max_dt
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the maximum creatinine level collected after the patient's arrival to the hospital.

Codes and Values:

- **Organ Dysfunction Hepatic Arrival/Max Datetimes** are situational because hospitals may not always have values to report
 - If any Organ Dysfunction Renal is reported then the datetime for Organ Dysfunction Renal value should be reported. For example, if Organ Dysfunction Renal Arrival has a value, Organ Dysfunction Renal Arrival Datetime should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)

hh = two digits of hour (00 through 23) (am/pm NOT allowed) mm = two digits of minute (00 through 59)

- 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
- 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:

Data Element Name: Template Variable: Format – Length: Mandatory:

Severity Variables

Organ Dysfunction Respiratory organ_dysfunc_respiratory Enumerated – 1 <mark>Yes</mark>

Description:

Indicates that the patient has respiratory organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No 1 = Yes

- Please see Appendix 5B for a list of applicable ICD-10-CM codes.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced appendix.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced appendix.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets 1
Template Variable:	platelets_1
Format – Length:	String — 6
Mandatory:	No

Indicates the first platelet level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets 2
Template Variable:	platelets_2
Format – Length:	String — 6
Mandatory:	No

Description:

Indicates the second platelet level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets 3
Template Variable:	platelets_3
Format – Length:	String — 6
Mandatory:	No

Description:

Indicates the third platelet level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets Min
Template Variable:	platelets_min
Format – Length:	String — 6
Mandatory:	No

Description:

Indicates the minimum platelet level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual platelet level. Convert the units to cells/uL if needed.

- *Platelets 1/2/3/Max* and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- Must be reported to one decimal place (example 2.8).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.
- This element belongs to Organ Dysfunction Hematologic.
- Formatting:
 - 1. Format must be a string up to 6-digits long.
 - 2. Example:
 - a. Platelet 320,000/uL should be reported as 320000
 - b. Platelet 60,000/uL should be reported as 60000

Dataset Segment:	Severity Variables
Data Element Name:	Platelets Datetime 1
Template Variable:	platelets_dt_1
Format – Length:	Datetime – 16
Mandatory:	No

Indicates the date and time of the first platelet level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets Datetime 2
Template Variable:	platelets_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second platelet level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets Datetime 3
Template Variable:	platelets_dt_3
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the third platelet level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Platelets Datetime Min
Template Variable:	<mark>platelets_dt_min</mark>
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	No

Indicates the date and time of the minimum platelet level collected after the patient's arrival to the hospital.

Codes and Values:

Notes for Abstraction:

- Platelets 1/2/3/Max Datetimes are situational because hospitals may not always have values to report.
 - If any Platelets are reported then the datetime for the Platelets value should be reported. For example, if *Platelets 1* has a value, *Platelets Datetime 1* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- This element belongs to Organ Dysfunction Hematologic.
- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - 2. YYYY = four-digit year

MM = two-digit month (01 = January, etc.)

- DD = two-digit day of month (01 through 31)
- hh = two digits of hour (00 through 23) (am/pm NOT allowed)

mm = two digits of minute (00 through 59)

- 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
- 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate 1
Template Variable:	sirs_heartrate_1
Format – Length:	Enumerated— 3
Mandatory:	Yes

Indicates the first heart rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate 2
Template Variable:	sirs_heartrate_2
Format – Length:	Enumerated — 3
Mandatory:	No

Description:

Indicates the second heart rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate 3
Template Variable:	sirs_heartrate_3
Format – Length:	Enumerated — 3
Mandatory:	No

Description:

Indicates the third heart rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate Max
Template Variable:	sirs_heartrate_max
Format – Length:	Enumerated — 3
Mandatory:	No

Description:

Indicates the maximum heart rate value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual heart rate.

- SIRS Heart Rate 1 and corresponding SIRS Heart Rate Datetime 1 are mandatory. However, SIRS Heart Rate 2, SIRS Heart Rate 3, and SIRS Heart Rate Max and corresponding datetimes are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - a. If *SIRS Heart Rate 2, SIRS Heart Rate 3,* and/or *SIRS Heart Rate Max* are collected then these values and their corresponding datetimes must be reported.
- Formatting:
 - 1. Format must be a number up to 3 digits.
 - 2. Example:
 - a. Heart rate/Pulse 100 beats per minutes (bpm) should be reported as 100
 - b. Heart rate/Pulse 43 beats per minutes (bpm) should be reported as 43

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate Datetime 1

Data Element Name: Template Variable: Format – Length: Mandatory:

SIRS Heart Rate Datetime 1 sirs heartrate dt 1

Datetime – 16

Yes

Description:

Indicates the date and time of the first heart rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate Datetime 2
Template Variable:	sirs_heartrate_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second heart rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate Datetime 3
Template Variable:	sirs_heartrate_dt_3
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the third heart rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Heart Rate Datetime Max
Template Variable:	sirs_heartrate_dt_max
Format – Length:	Datetime – 16
Mandatory:	No No

Indicates the date and time of the maximum heart rate value collected after the patient's arrival to the hospital.

Codes and Values:

- SIRS Heart Rate 1 Datetime is mandatory. However, SIRS Heart Rate Datetime 2, SIRS Heart Rate Datetime 3, SIRS Heart Rate Datetime Max are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - If any Heart Rate is reported then the datetime for the Heart Rate value should be reported. For example, if SIRS Heart Rate 2 has a value, then SIRS Heart Rate Datetime 2 should not be blank.
- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Leukocyte Arrival
Template Variable:	sirs_leuckocyte_arrival
Format – Length:	String — 6

No

Description:

Mandatory:

Indicates the first white blood cell (WBC) level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Leukocyte Min
Template Variable:	sirs_leuckocyte_min
Format – Length:	String — 6
Mandatory:	No

Description:

Indicates the minimum white blood cell (WBC) level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Leukocyte Max
Template Variable:	sirs_leuckocyte_max
Format – Length:	String — 6
Mandatory:	No

Description:

Indicates the maximum white blood cell (WBC) level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual white blood cell (WBC) count. Convert the units to cells/uL if needed.

- *SIRS Leukocyte Arrival/Min/Max* and corresponding datetimes are situational because hospitals may not always have values to report.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.

- Must be reported to one decimal place (example 2.8).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example <0.1, should be reported as <0.1.

<u>Formatting:</u>

- 1. Format must be a string up to 6-digits.
- 2. Example:
 - WBC 100,000/uL should be reported as 100000
 - WBC 11,500/uL should be reported as 11500
 - o WBC 4,400/uL should be reported as 4400

Dataset Segment:

Data Element Name: Template Variable: Format – Length: Mandatory:

Severity Variables

SIRS Leukocyte Arrival Datetime sirs_leuckocyte_arrival_dt Datetime – 16 No

Description

Indicates the date and time of the first white blood cell (WBC) collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Leukocyte Min Datetime
Template Variable:	sirs_leuckocyte_min_dt
Format – Length:	Datetime – 16
Mandatory:	No

Description

Indicates the date and time of minimum white blood cell (WBC) level collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Leukocyte Max Datetime
Template Variable:	sirs leuckocyte max dt
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	<mark>No</mark>

Description

Indicates the date and time of maximum white blood cell (WBC) level collected after the patient's arrival to the hospital.

Codes and Values:

Notes for Abstraction:

• SIRS Leukocyte Arrival/Min/Max Datetimes are situational because hospitals may not always have values to report.

- If any SIRS Leukocyte is reported then the datetime for the SIRS Leukocyte value should be reported. For example, if *SIRS Leukocyte Arrival* has a value, *SIRS Leukocyte Arrival Datetime* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - 2. YYYY = four-digit year
 - MM = two-digit month (01 = January, etc.) DD = two-digit day of month (01 through 31) hh = two digits of hour (00 through 23) (am/pm NOT allowed) mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate 1
Template Variable:	sirs_respiratoryrate_1

Number — 2

Yes

Format – Length: <mark>Mandatory:</mark>

Description:

Indicates the first respiratory rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate 2
Template Variable:	sirs_respiratoryrate_2
Format – Length:	Number — 2
Mandatory:	No

Description:

Indicates the second respiratory rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate 3
Template Variable:	sirs_respiratoryrate_3
Format – Length:	Number — 2
Mandatory:	No

Description:

Indicates the third respiratory rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate Max
Template Variable:	sirs_respiratoryrate_max
Format – Length:	Number — 2
Mandatory:	No

Description:

Indicates the maximum respiratory rate value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual respiratory rate.

- SIRS Respiratory Rate 1 and corresponding SIRS Respiratory Rate Datetime 1 are mandatory. However, SIRS Respiratory Rate 2, SIRS Respiratory Rate 3, and SIRS Respiratory Rate Max and corresponding datetimes are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - a. If SIRS Respiratory Rate 2, SIRS Respiratory Rate 3, and/or SIRS Respiratory Rate Max are collected then these values and their corresponding datetimes must be reported.
- Formatting:
 - 1. Format must be a number up to 2-digits.
 - 2. Example:
 - a. Respiratory rate 12 breaths per minutes (bpm) should be reported as 12
 - b. Respiratory rate 9 breaths per minutes (bpm) should be reported as 9

Dataset Segment:

Data Element Name: Template Variable: Format – Length: Mandatory:

Severity Variables

SIRS Respiratory Rate Datetime 1 sirs_respiratoryrate_dt_1 Datetime – 16 Yes

Description:

Indicates the date and time of the first respiratory rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate Datetime 2
Template Variable:	sirs_respiratoryrate_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second respiratory rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate Datetime 3
Template Variable:	sirs_respiratoryrate_dt_3
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the third respiratory rate value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Respiratory Rate Datetime Max
Template Variable:	sirs_respiratoryrate_dt_max
Format – Length:	Datetime – 16
Mandatory:	No

Indicates the date and time of the maximum respiratory rate value collected after the patient's arrival to the hospital.

Codes and Values:

- SIRS Respiratory Rate 1 Datetime is mandatory. However, SIRS Respiratory Rate Datetime
 2, SIRS Respiratory Rate Datetime 3, SIRS Respiratory Rate Datetime Max are situational
 because hospitals may not always have values to report (i.e., additional values not collected).
 - If any SIRS Respiratory Rate is reported then the corresponding datetime should be reported. For example, if SIRS Respiratory Rate 2 has a value, then SIRS Respiratory Rate Datetime 2 should not be blank.
- Formatting:
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature 1
Template Variable:	sirs_temperature_1
Format – Length:	Enumerated — 5
Mandatory:	Yes

Indicates the first temperature value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature 2
Template Variable:	sirs_temperature_2
Format – Length:	Enumerated — 5
Mandatory:	No

Description:

Indicates the second temperature value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature 3
Template Variable:	sirs_temperature_3
Format – Length:	Enumerated — 5
Mandatory:	No

Description:

Indicates the third temperature value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature Max
Template Variable:	sirs_temperature_max
Format – Length:	Enumerated — 5
Mandatory:	No

Description:

Indicates the maximum temperature value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual temperature level using Fahrenheit. Convert from Celsius if needed.

- SIRS Temperature 1 and corresponding SIRS Temperature Datetime 1 are mandatory. However, SIRS Temperature 2, SIRS Temperature 3, and SIRS Temperature Max and corresponding datetimes are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - a. If *SIRS Temperature 2, SIRS Temperature 3,* and/or *SIRS Temperature Max* are collected then these values and their corresponding datetimes should be reported.
- Formatting:
 - 1. Must be numeric to one decimal place (example 98.8)
 - 2. Example:
 - a. 100.4°F should be reported as 100.4
 - b. 96°F should be reported as 96.0
 - c. 97.6°F should be reported as 97.6

Dataset Segment:	Severity Variables

Data Element Name: Template Variable: Format – Length: Mandatory:

SIRS Temperature Datetime 1 sirs_temperature_dt_1 Datetime – 16 Yes

Description:

Indicates the date and time of the first temperature value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature Datetime 2
Template Variable:	sirs_temperature_dt_2
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second temperature value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature Datetime 3
Template Variable:	sirs_temperature_dt_3
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the third temperature value collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	SIRS Temperature Datetime Max
Template Variable:	sirs_temperature_dt_max
Format – Length:	Datetime – 16
Mandatory:	No

Indicates the date and time of the maximum temperature value collected after the patient's arrival to the hospital.

Codes and Values:

Notes for Abstraction:

- SIRS Temperature Datetime 1 is mandatory. However, SIRS Temperature Datetime 2, SIRS Temperature Datetime 3, SIRS Temperature Datetime Max are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - If any Temperature is reported then the datetime for the Temperature value should be reported. For example, if SIRS Temperature 2 has a value, then SIRS Temperature Datetime 2 should not be blank.

Formatting:

- 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
- YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
- 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
- 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	Systolic First
Template Variable:	systolic_1
Format – Length:	Number – 3
Mandatory:	Yes

Indicates the patient's first systolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Second
Template Variable:	systolic_2
Format – Length:	Number – 3
Mandatory:	No

Description:

Indicate the patient's second systolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Third
Template Variable:	systolic_3
Format – Length:	Number – 3
Mandatory:	No

Description:

Indicate the patient's third systolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Min
Template Variable:	systolic_min
Format – Length:	Number – 3
<mark>Mandatory:</mark>	No

Indicates the patient's minimum systolic blood pressure collected after the patient's arrival to the hospital.

Codes and Values:

- Systolic First and corresponding Systolic First Datetime 1 are mandatory. However, Systolic Second, Systolic Third, and Systolic Min and corresponding datetimes are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - If any Systolic is reported then the datetime for the Systolic value should be reported. For example, if *Systolic Second* has a value, then *Systolic Second Datetime 2* should not be blank.
- Formatting:
- Format must be a number up to 3 digits.
 - 2. Example:
 - a. Systolic blood pressure 80mm Hg should be reported as 80
 - b. Systolic blood pressure 112 Hg should be reported as 112

Dataset Segment:Severity VariablesData Element Name:Systolic First Datetime 1

Data Element Name: Template Variable: Format – Length: <mark>Mandatory:</mark> Systolic First Datetime 1 systolic_dt_1 Datetime – 16 Yes

Description:

Indicates the date and time of the first systolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Second Datetime 2
Template Variable:	<mark>systolic _dt_2</mark>
Format – Length:	Datetime – 16
Mandatory:	No

Description:

Indicates the date and time of the second systolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Third Datetime 3
Template Variable:	<mark>systolic_dt_3</mark>
Format – Length:	Datetime – 16
<mark>Mandatory:</mark>	No

Description:

Indicates the date and time of the third systolic blood pressure collected after the patient's arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Datetime Min
Template Variable:	<mark>systolic_dt_min</mark>
Format – Length:	Datetime – 16
Mandatory:	No

Indicates the date and time of the minimum systolic blood pressure collected after the patient's arrival to the hospital.

Codes and Values:

- Systolic First Datetime 1 is mandatory. However, Systolic Second Datetime 2, Systolic Third Datetime 3, Systolic Datetime Min are situational because hospitals may not always have values to report (i.e., additional values not collected).
 - If any Systolic is reported then the datetime for the Systolic value should be reported. For example, if *Systolic Second* has a value, then *Systolic Second Datetime 2* should not be blank.
- <u>Formatting:</u>
 - 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 - YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 - 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 a. 1959-11-03T23:42 is also valid
 - 4. Midnight = 00:00, **NOT** 24:00

Change Log

Version D2.1

- Please review *Key points to remember during data extraction* in detail as several paragraphs were modified.
- One code (J1282 Pneumonia due to coronavirus disease 2019) was added to Table B in *Inclusion Definition* on page 10.
- One code (R0602 Shortness of breath) on Table B in *Inclusion Definition* on page 11 was corrected to include a missing 0.
- *Ethnicity* was updated to align with SPARCS code set.
- Data Element Name of History of COVID-19 Date changed to History of COVID-19 Datetime
- Five new data elements were added:
 - Other Payer
 - o Transferred In
 - o Transferred Out
 - Transfer Facility Name Receiving
 - Transfer Facility Name Sending
- Description was updated for the following variables:
 - Transfer Facility Identifier Receiving
 - Transfer Facility Identifier Sending
- The Data Segment section *Header* was changed to include the data element reporting requirement to indicate mandatory yes or no. The following variables were changed from mandatory to optional because hospitals may not always have values to report.
 - History of COVID-19 Datetime
 - Patient Care Considerations Date
 - o **aPTT 1**
 - o **aPTT 2**
 - o **aPTT 3**
 - o aPTT Max
 - o aPTT Datetime 1
 - o aPTT Datetime 2
 - o aPTT Datetime 3
 - o aPTT Datetime Max
 - Arrival Datetime
 - o Diastolic Second
 - o Diastolic Third
 - o Diastolic Min
 - Diastolic Second Datetime 2
 - Diastolic Third Datetime 3
 - Diastolic Datetime Min
 - o **INR 1**
 - o INR 2

- o INR 3
- o INR Max
- o INR Datetime 1
- o INR Datetime 2
- o INR Datetime 3
- INR Datetime Max
- o Lactate Level 1
- Lactate Level 2
- Lactate Level 3
- o Lactate Level Max
- Lactate Level Datetime 1
- o Lactate Level Datetime 2
- Lactate Level Datetime 3
- o Lactate Level Datetime Max
- Organ Dysfunction Hepatic Arrival
- Organ Dysfunction Hepatic Max
- o Organ Dysfunction Hepatic Arrival Datetime
- Organ Dysfunction Hepatic Max Datetime
- Organ Dysfunction Renal Arrival
- Organ Dysfunction Renal Max
- o Organ Dysfunction Renal Arrival Datetime
- o Organ Dysfunction Renal Max Datetime
- o Platelets 1
- o Platelets 2
- o Platelets 3
- Platelets Min
- o Platelets Datetime 1
- o Platelets Datetime 2
- o Platelets Datetime 3
- Platelets Datetime Min
- SIRS Heart Rate 2
- o SIRS Heart Rate 3
- SIRS Heart Rate Max
- SIRS Heart Rate Datetime 2
- SIRS Heart Rate Datetime 3
- o SIRS Heart Rate Datetime Max
- SIRS Leukocyte Arrival
- SIRS Leukocyte Min
- o SIRS Leukocyte Max
- SIRS Leukocyte Arrival Datetime
- SIRS Leukocyte Min Datetime
- SIRS Leukocyte Max Datetime
- o SIRS Respiratory Rate 2
- SIRS Respiratory Rate 3

- SIRS Respiratory Rate Max
- SIRS Respiratory Rate Datetime 2
- SIRS Respiratory Rate Datetime 3
- SIRS Respiratory Rate Datetime Max
- SIRS Temperature 2
- o SIRS Temperature 3
- SIRS Temperature Max
- SIRS Temperature Datetime 2
- SIRS Temperature Datetime 3
- o SIRS Temperature Datetime Max
- Systolic Second
- Systolic Third
- Systolic Min
- Systolic Second Datetime 2
- Systolic Third Datetime 3
- o Systolic Datetime Min
- Format Length, Notes of Abstraction were updated for the following variables:
 - ICD-10-CM Code (n)
 - o **Payer**
 - History of Other Cardiovascular Disease
 - o **Race**
 - Patient Care Considerations
 - Suspected Source of Infection
 - o **aPTT 1**
 - o **aPTT 2**
 - o **aPTT 3**
 - o aPTT Max
 - o aPTT Datetime 1
 - o aPTT Datetime 2
 - o aPTT Datetime 3
 - o aPTT Datetime Max
 - Diastolic First
 - Diastolic Second
 - o Diastolic Third
 - Diastolic Min
 - Distolic First Datetime 1
 - Diastolic Second Datetime 2
 - o Diastolic Third Datetime 3
 - o Diastolic Datetime Min
 - o **INR 1**
 - o INR 2
 - o INR 3
 - o INR Max
 - o INR Datetime 1

- o INR Datetime 2
- o INR Datetime 3
- o INR Datetime Max
- o Lactate Level 1
- o Lactate Level 2
- o Lactate Level 3
- o Lactate Level Max
- Lactate Level Datetime 1
- o Lactate Level Datetime 2
- o Lactate Level Datetime 3
- Lactate Level Datetime Max
- Organ Dysfunction CNS
- Organ Dysfunction Hepatic Arrival
- Organ Dysfunction Hepatic Max
- o Organ Dysfunction Hepatic Arrival Datetime
- Organ Dysfunction Hepatic Max Datetime
- Organ Dysfunction Renal Arrival
- Organ Dysfunction Renal Max
- o Organ Dysfunction Renal Arrival Datetime
- Organ Dysfunction Renal Max Datetime
- Organ Dysfunction Respiratory
- o Platelets 1
- o Platelets 2
- o Platelets 3
- o Platelets Min
- o Platelets Datetime 1
- Platelets Datetime 2
- Platelets Datetime 3
- Platelets Datetime Min
- o SIRS Heart Rate 1
- o SIRS Heart Rate 2
- SIRS Heart Rate 3
- SIRS Heart Rate Max
- o SIRS Heart Rate Datetime 1
- SIRS Heart Rate Datetime 2
- SIRS Heart Rate Datetime 3
- o SIRS Heart Rate Datetime Max
- SIRS Leukocyte Arrival
- SIRS Leukocyte Min
- SIRS Leukocyte Max
- SIRS Leukocyte Arrival Datetime
- SIRS Leukocyte Min Datetime
- SIRS Leukocyte Max Datetime
- SIRS Respiratory Rate 1

- SIRS Respiratory Rate 2
- SIRS Respiratory Rate 3
- SIRS Respiratory Rate Max
- SIRS Respiratory Rate Datetime 1
- SIRS Respiratory Rate Datetime 2
- SIRS Respiratory Rate Datetime 3
- SIRS Respiratory Rate Datetime Max
- SIRS Temperature 1
- o SIRS Temperature 2
- o SIRS Temperature 3
- SIRS Temperature Max
- o SIRS Temperature Datetime 1
- SIRS Temperature Datetime 2
- SIRS Temperature Datetime 3
- SIRS Temperature Datetime Max
- Systolic Second
- Systolic Third
- Systolic Min
- Systolic Second Datetime 2
- Systolic Third Datetime 3
- Systolic Datetime Min
- Variable names were changed for the following variables from "datetime" to "dt" to provide more consistency:
 - Admission Datetime
 - Arrival Datetime
 - o Discharge Datetime
 - o aPTT Datetime 1
 - o aPTT Datetime 2
 - o aPTT Datetime 3
 - o aPTT Datetime Max
 - Diastolic First Datetime 1
 - Diastolic Second Datetime 2
 - o Diastolic Third Datetime 3
 - Diastolic Datetime Min
 - o INR Datetime 1
 - o INR Datetime 2
 - o INR Datetime 3
 - INR Datetime Max
 - o Lactate Level Datetime 1
 - Lactate Level Datetime 2
 - o Lactate Level Datetime 3
 - Lactate Level Datetime Max
 - o Platelets Datetime 1
 - Platelets Datetime 2

- Platelets Datetime 3
- Platelets Datetime Min
- Systolic First Datetime 1
- Systolic Second Datetime 2
- Systolic Third Datetime 3
- Systolic Datetime Min
- Several appendices were modified to remove decimal places within codes. Be sure to review Appendices D2.1 change log for the specific variables that were updated. The following appendices were modified:
 - Appendix 1Y: Smoking or Vaping Comorbidity / Risk Factor (POA) ICD-10 CM Codes
 - o Appendix 2D: Flu Positive Clinical ICD-10-CM Codes
- Appendix 2A: COVID-19 Exposure Clinical ICD-10-CM Codes was updated to include one additional code.
- Appendix 2E: Suspected Source of Infection was updated to include additional codes.
- Be sure to review Appendices D2.1 for the specific codes that were added.

Version D2.0

- *Suspected Source of Infection* value was changed for option 11 to read "soft tissue infection".
- The order of the appendices was changed to align with the order of the data elements in the data dictionary.
- Several appendices were updated to include additional codes. Be sure to review Appendices D2.0 for the specific codes that were added. The following appendices were modified:
 - Appendix 1E: Chronic Liver Disease Comorbidity / Risk Factor
 - o Appendix 1G: Chronic Respiratory Failure Comorbidity / Risk Factor
 - o Appendix 1H: Coagulopathy Comorbidity / Risk Factor
 - o Appendix 1P: Immunocompromising Comorbidity / Risk Factor
 - Appendix 1S: Medication Anticoagulant
 - Appendix 1T: Medication Immune Modifying
 - Appendix 1V: Obesity Comorbidity
 - Appendix 2D: Flu Positive Clinical
 - Appendix 2E: Suspected Source of Infection
 - Appendix 3F: Vasopressor Administration Treatment Medication and NDC Codes
- Descriptions of CSV files of codes were added to *Key points to remember during data extraction*.
- "Collected" was added to the severity variables to specify that collected datetime should be reported for the labs and vital signs.
- Cardiac outcome was renamed to cardiovascular outcome.

- The notes for abstraction of *Race* were modified to provide a single link to SPARCS codes and values.
- The notes for abstraction of *Pregnancy Status During Hospitalization* were modified.
- The notes for abstraction of *Flu Positive* were modified.
- The notes for abstraction of *Tracheostomy in Hospital* were modified.
- The description of *Medication Immune Modifying* was modified.
- The description and codes and values of *Lactate Level Max* were modified.
- The notes for abstraction of all labs reported as Severity Variables have been modified. The following notes have been added for the labs and their respective collected date and time:
 - For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
 - For all labs, if the numeric value is preceded by "<" (less than) or ">" (greater than) signs, this sign should be reported. For example <0.1 should be reported as <0.1.
 For example >100,000 should be reported as >100,000.