

NYSDOH Pediatric Sepsis and COVID-19 Data Dictionary

Digitalized Data Collection, D2.0

Version (Digital) D2.0

February 10, 2022

**Effective with
Discharges from 01/01/2022**

This dictionary refers to administrative codes available for download as csv files to assist in data extraction.

The most recent version of this document, the *Frequently Asked Questions* document, the *Table of Elements* data template, and the instructions may be found at:

<https://ny.sepsis.ipro.org>

Questions regarding this document should be submitted at:

<https://ny.sepsis.ipro.org/support>

Changes from version D1.1 to D2.0 are highlighted in yellow.

Prepared by:
Office of Quality and Patient Safety
New York State Department of Health
Revised February 10, 2022

Table of Contents

Key points to remember during data extraction	8
Inclusion Definition.....	10
Demographic Variables.....	12
Admission Datetime	13
Arrival Datetime.....	14
Date of Birth.....	16
Discharge Datetime	17
Discharge Status	18
Ethnicity	20
Facility Identifier	21
Gender	22
ICD-10-CM Code (n).....	23
ICD-10-CM POA Indicator (n).....	24
Inclusion COVID-19	26
Inclusion Septic Shock.....	27
Inclusion Severe Sepsis	28
Insurance Number	29
Medical Record Number.....	30
Other Payer.....	31
Patient Control Number	32
Patient City.....	33
Patient County Code	34
Patient Street Address Line 1	35
Patient Street Address Line 2	36
Patient State	37
Patient Zip Code of Residence.....	38
Payer	39
Race.....	40
Source of Admission	41
Transferred In	42
Transferred Out	43

Transfer Facility Identifier Receiving	44
Transfer Facility Identifier Sending	45
Transfer Facility Name Receiving	46
Transfer Facility Name Sending	47
Unique Personal Identifier	48
Comorbidity/Risk Factor Variables	50
Acute Cardiovascular Conditions (POA)	51
Acquired Absent Spleen	53
Adrenal Gland Disorder (POA)	54
AIDS/HIV Disease	56
Asthma	57
Bronchopulmonary Dysplasia	58
Chronic Liver Disease	59
Chronic Kidney Disease	60
Chronic Respiratory Failure	61
Coagulopathy (POA)	62
Congestive Heart Failure	64
Cystic Fibrosis	65
Diabetes	66
Dialysis Comorbidity (POA)	67
History of COVID -19	69
History of COVID-19 Datetime	70
History of Other Cardiovascular Disease	71
Hypertension	72
Immunocompromising	73
Kawasaki/Mucocutaneous Lymph Node Syndrome	74
Lymphoma Leukemia Multiple Myeloma	75
Mechanical Ventilation Comorbidity (POA)	76
Medication Immune Modifying Pre-Hospital	78
Metastatic Cancer	79
Neutropenic Patients (POA)	80
Obesity	82
Patient Care Considerations	83

Patient Care Considerations Date.....	84
Pregnancy Comorbidity	85
Pregnancy Status During Hospitalization	86
Prematurity	87
Sickle Cell Disease	88
Skin Disorders/Burns	89
Smoking Vaping	90
Stem Cell Transplant Recipients	91
Clinical Variables	92
COVID-19 Exposure.....	93
COVID-19 Virus	94
Drug Resistant Pathogen	95
Flu Positive.....	96
Suspected Source of Infection	97
Treatment (in hospital) Variables	98
Dialysis Treatment	99
During Hospital Immune Modifying Medication.....	100
During Hospital Remdesivir	101
ECMO	102
High Flow Nasal Cannula	103
IVIg.....	104
Mechanical Ventilation Treatment.....	105
Non-Invasive Positive Pressure Ventilation.....	106
Vasopressor Administration	107
Outcome (at discharge) Variables	108
Dialysis Outcome	109
Mechanical Ventilation Outcome.....	110
Neurological Outcome.....	111
Outcome (in hospital) Variables	112
Cardiovascular Outcomes in Hospital.....	113
ICU During Hospitalization.....	114
Severity Variables	115
aPTT 1.....	116

aPTT 2.....	116
aPTT 3.....	116
aPTT Max	116
aPTT Datetime 1	118
aPTT Datetime 2	118
aPTT Datetime 3	118
aPTT Datetime Max	118
Bilirubin Arrival	120
Bilirubin Max.....	120
Bilirubin Arrival Datetime	122
Bilirubin Max Datetime.....	122
Creatinine Arrival	124
Creatinine Max.....	124
Creatinine Arrival Datetime.....	126
Creatinine Max Datetime	126
Diastolic First.....	128
Diastolic Second.....	128
Diastolic Third	128
Diastolic Min	128
Diastolic First Datetime 1.....	130
Diastolic Second Datetime 2.....	130
Diastolic Third Datetime 3	130
Diastolic Datetime Min	130
INR 1.....	132
INR 2.....	132
INR 3.....	132
INR Max.....	132
INR Datetime 1.....	134
INR Datetime 2.....	134
INR Datetime 3.....	134
INR Datetime Max.....	134
Lactate Level 1	136
Lactate Level 2	136

Lactate Level 3	136
Lactate Level Max	136
Lactate Level Datetime 1	138
Lactate Level Datetime 2	138
Lactate Level Datetime 3	138
Lactate Level Datetime Max	138
Organ Dysfunction Cardiovascular	140
Organ Dysfunction CNS.....	141
Organ Dysfunction Hematologic.....	142
Organ Dysfunction Hepatic.....	143
Organ Dysfunction Renal	144
Organ Dysfunction Respiratory	145
Platelets 1	146
Platelets 2	146
Platelets 3	146
Platelets Min	146
Platelets Datetime 1	148
Platelets Datetime 2	148
Platelets Datetime 3	148
Platelets Datetime Min	148
SIRS Heart Rate 1	150
SIRS Heart Rate 2	150
SIRS Heart Rate 2	150
SIRS Heart Rate Max	150
SIRS Heart Rate Datetime 1	152
SIRS Heart Rate Datetime 2	152
SIRS Heart Rate Datetime 3	152
SIRS Heart Rate Datetime Max	152
SIRS Leukocyte Arrival	154
SIRS Leukocyte Min.....	154
SIRS Leukocyte Max	154
SIRS Leukocyte Arrival Datetime	156
SIRS Leukocyte Min Datetime.....	156

SIRS Leukocyte Max Datetime	156
SIRS Respiratory Rate 1.....	158
SIRS Respiratory Rate 2.....	158
SIRS Respiratory Rate 3.....	158
SIRS Respiratory Rate Max.....	158
SIRS Respiratory Rate Datetime 1.....	160
SIRS Respiratory Rate Datetime 2.....	160
SIRS Respiratory Rate Datetime 3.....	160
SIRS Respiratory Rate Datetime Max	160
SIRS Temperature 1	162
SIRS Temperature 2	162
SIRS Temperature 3	162
SIRS Temperature Max	162
SIRS Temperature Datetime 1	164
SIRS Temperature Datetime 2	164
SIRS Temperature Datetime 3	164
SIRS Temperature Datetime Max	164
Systolic First	166
Systolic Second	166
Systolic Third.....	166
Systolic Min.....	166
Systolic First Datetime 1	168
Systolic Second Datetime 2	168
Systolic Third Datetime 3.....	168
Systolic Datetime Min.....	168
Change Log	170

Key points to remember during data extraction

The New York State Department of Health (NYSDOH) is collecting data for all inpatient, emergency department (ED), and observation pediatric patients with a diagnosis of severe sepsis and septic shock, as defined by the case inclusion definition on pages 10-11 of this data dictionary. NYSDOH is collecting data for all inpatient pediatric patients with a COVID-19 or MIS-C diagnosis as defined by the case inclusion definition on pages 10-11 of this data dictionary. Please note that patients with a COVID-19 or MIS-C diagnosis but no severe sepsis or septic shock diagnosis that are evaluated in the emergency department or under hospital observation status but are never admitted for inpatient care should **not** be reported.

Data for all patients under 21 years of age at the time of admission and meet the case *inclusion definition*, are to be reported into the pediatric NYSDOH database. For patients who are admitted to the hospital, the patients' age at admission should be used to determine reporting to the adult or the pediatric database. For patients who are not admitted to the hospital (i.e., ED/Observation only), the patients' age at arrival should be used to determine reporting to the adult or the pediatric database.

Because the information in the appendices and csv file is duplicative, appendices will no longer be made available beginning with Data Dictionary Version D2.0. Any references to the appendices have been replaced with references to the corresponding csv files. When using the csv files for the identification of relevant ICD-10-CM codes, be sure to capture any code (ICD-10-CM) in any position at any point during hospitalization unless otherwise indicated in the variable directions.

Most variables are required but there are some exceptions such as *Transfer Facility Identifier Receiving* and *Transfer Facility Identifier Sending* which are situational. For transfer data elements we recognize that your hospital EHR may not have Transfer Facility Identifier Receiving/Sending but may have Transfer Facility Name Receiving/Sending. Please report all data you have regarding transfers.

Hospitals that have within hospital transfer patients (i.e., patient transferred from one unit to another within the same hospital) should report the case as it is collected in the EHR. For example, if your EHR represents a patient transferred from a rehab unit to an acute care unit as a combined record in your EHR, please report this episode of care as one record, even if two separate bills are generated for the rehab and the inpatient admission. If there are two separate records in the EHR, please submit it as two separate cases if inclusion criteria are met for each case. Be sure to use the appropriate discharge disposition to accurately represent the case.

This data dictionary has been designed to eliminate the need for manual chart abstraction and to permit hospitals to utilize their information technology staff and electronic medical record systems to extract the necessary data. This data will be accepted into the current portal in a flat file format following existing procedures which may be found at <https://ny.sepsis.ipro.org/>.

The csv files of codes are provided separately. Each csv file contains three columns: the codes for the variable, the corresponding code description, and a subcategory if applicable. For example:

ICD-10-CM CODE	ICD-10-CM CODE DESCRIPTION	Subcategory
I2101	ST elevation (STEMI) myocardial infarction involving left main coronary artery	MI
I2102	ST elevation (STEMI) myocardial infarction involving left anterior descending coronary artery	MI

Please note that data elements with multiple selections (more than Yes/No) will have values/contents in the subcategory column in the csv files, for example *Acute Cardiovascular Conditions*. In general, the naming convention for csv files is TemplateVariable_code_Version. For example:

- ped_asthma_code_VerD2.0.csv

In the event that the csv files are for NDC codes of medications, ndc is added in the naming convention. For example:

- ped_medication_immune_modifying_ndc_code_VerD2.0.csv

Inclusion Definition

For all cases of severe sepsis and/or septic shock with or without COVID-19/MIS-C, the pediatric database includes inpatient, ED and Observation cases. For COVID-19/MIS-C cases without severe sepsis and/or septic shock, the pediatric database is limited to inpatient cases only. The pediatric database also includes all newborn patients, including those who have expired during birth hospitalization. For patients who are admitted to the hospital, the patients' age at admission should be used to determine reporting to the adult or the pediatric database. For patients who are not admitted to the hospital (i.e., ED and Observation only), the patients' age at arrival should be used to determine reporting to the adult or the pediatric database. Hospitals should include admissions as defined by the ICD-10-CM codes delineated below.

The NYSDOH is identifying the (denominator) population of cases for inclusion into the database using ICD-10-CM codes. Hospitals may use all sources of data for case inclusion (electronic medical record codes as well as administrative and billing codes); however, cases should only be reported if one of the below inclusion codes is a final diagnosis. This will allow for electronic identification of cases. The ICD-10-CM code-based definition for identifying the severe sepsis/septic shock and COVID-19/MIS-C patient population for abstraction includes the following codes which are presented in Tables A and B. Cases with codes in either table are to be reported.

Hospitals will report cases where criteria are met by:

- At least one code in Table A; OR
- At least one code in Table B

Examples:

- Patient with Code T8112XA and no other code from Table A or Table B is reported.
- Patient with Code U072 and no other code from Table A or Table B is reported.

Table A: Severe sepsis and/or septic shock inclusion ICD-10-CM codes

Severe Sepsis/Septic Shock	
ICD-10-CM	Description
R6520	Severe sepsis without septic shock
R6521	Severe sepsis with septic shock
T8112XA	Post procedural septic shock, initial encounter

OR

Table B: COVID-19 inclusion ICD-10-CM codes

COVID-19		
ICD-10-CM	Description	Type
U071	COVID-19, virus identified	COVID-19

COVID-19		
ICD-10-CM	Description	Type
U072	COVID-19, virus not identified (Clinically-epidemiologically diagnosed COVID-19)	COVID-19
J1282	Pneumonia due to coronavirus disease 2019 *The code J1282 is effective as of January 1, 2021	COVID-19
M3581	Multisystem Inflammatory syndrome (MIS) *The code M3581 is effective as of January 1, 2021	MIS-C

Demographic Variables

Dataset Segment:**Demographic Variables**

Data Element Name:

Admission Datetime

Template Variable:

admission_dt

Format – Length:

Datetime – 16

Mandatory:

Yes

Description:

Indicates the date and time that the patient was admitted to inpatient status at the hospital.

Codes and Values:

Enter the Admission Datetime.

Notes for Abstraction:

- **Formatting:**
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm is **NOT** allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00
- Cannot have been after *Discharge Datetime*.
- If there is a difference between arrival to inpatient floor and the written admission order, report the time the admission order was written.

Dataset Segment:**Demographic Variables**

Data Element Name:

Arrival Datetime

Template Variable:

arrival_dt

Format – Length:

Datetime – 16

Mandatory:

Yes

Description:

Indicates the earliest documented date and time the patient arrived at the hospital.

Codes and Values:

Enter the Arrival Datetime.

Notes for Abstraction:

- **Formatting:**
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm is **NOT** allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00
- Report earliest date and time the patient arrived at the ED, at the nursing floor, for observation, or as a direct admit to the cath lab.
- The arrival date and time may differ from the *Admission Datetime*.
- Cannot be after the *Discharge Datetime*.
- **Observation Status: Only include these observation cases if admitted to inpatient status.**
 - If the patient was admitted to observation from an outpatient setting of the hospital, use the date and time the patient arrives at the ED or on the floor of observation care as the arrival date and time.
 - If the patient was admitted to observation from the ED of the hospital, use the date and time the patient arrived at the ED as the *Arrival Datetime*.
- **Direct Admits:**
 - If the patient is a “Direct Admit” to the cath lab, use the earliest date and time the patient arrived at the cath lab (or cath lab staging/holding area) as the *Arrival Datetime*.

- If the patient is a “Direct Admit” to acute inpatient or observation, use the earliest date and time the patient arrived at the nursing floor or in observation as the *Arrival Datetime*.
- If the patient was transferred from your hospital’s satellite/free-standing ED or from another hospital within your hospital’s system (as an inpatient or ED patient) and there is one medical record for the care provided at both facilities, use the *Arrival Datetime* at the first facility.
- The *Arrival Datetime* can be obtained from the time period that the patient was an ED patient.

Dataset Segment:**Demographic Variables**

Data Element Name:

Date of Birth

Template Variable:

date_of_birth

Format – Length:

Date — 10

Mandatory:

Yes

Description:

Indicates the date of birth of the patient.

Codes and Values:

Enter the Date of Birth.

Notes for Abstraction:

- **Formatting:**
 1. Format must be YYYY-MM-DD
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
 3. Example: November 3, 1959 = 1959-11-03
- ***Date of Birth*** cannot be after ***Admission Datetime***.
- Patient age at admission should be used to determine reporting to the adult or the pediatric database.
- Data for all patients who are under 21 years of age are to be reported into the pediatric NYSDOH database.
 - Patients 21 years of age or older as of their admission date will be rejected and required for submission to the adult sepsis data file.

Dataset Segment:**Demographic Variables**

Data Element Name:

Discharge Datetime

Template Variable:

discharge_dt

Format – Length:

Datetime — 16

Mandatory:

Yes

Description:

Indicates the date and time that the patient was discharged from the hospital, left against medical advice, or expired.

Codes and Values:

Enter the Discharge Datetime.

Notes for Abstraction:

- **Formatting:**
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm **NOT** allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00
- Cannot precede 2014-04-01 00:00.
- Cannot precede *Admission Datetime* or *Arrival Datetime*.
- If the time of death and administrative discharge date and times are not the same, use the time of death for *Discharge Datetime*.
- For a patient who is discharged from one unit/department to another unit/department within the same facility, the **final discharge from the facility** is what should be reported for *Discharge Datetime*. Do not use discharges from internal transfers, since these are not actually separate hospital admissions – the entire period should be submitted as one record.

Dataset Segment:**Demographic Variables**

Data Element Name:	Discharge Status
Template Variable:	discharge_status
Format – Length:	Enumerated – 2
Mandatory:	Yes

Description:

Indicates the code that best represents the patient’s destination after discharge from the hospital.

Codes and Values:

- 01 = Discharge to Home or Self Care (Routine Discharge). Includes discharge to home; home on oxygen if DME only; any other DME only; group home, foster care, independent living and other residential care arrangements; outpatient programs, such as partial hospitalization or outpatient chemical dependency programs.
- 02 = Discharged/transferred to a Short-Term General Hospital for Inpatient Care.
- 03 = Discharged/transferred to Skilled Nursing Facility (SNF) with Medicare Certification in anticipation of Skilled Care. Medicare indicates that the patient is discharged/transferred to a Medicare certified nursing facility. For hospitals with an approved swing bed arrangement, use Code 61 Swing Bed. For reporting other discharges/transfers to nursing facilities see 04 and 64.
- 04 = Discharged/transferred to a Facility that Provides Custodial or Supportive Care. This is used to designate patients that are discharged/transferred to a nursing facility with neither Medicare nor Medicaid certification and for discharges/transfers to Assisted Living Facilities.
- 05 = Discharged/transferred to a Designated Cancer Center or Children's Hospital.
- 06 = Discharged/transferred to Home under Care of Organized Home Health Service Organization in Anticipation of Covered Skilled Care. Report this code when the patient is discharged/transferred to home with a written plan of care (tailored to the patient's medical needs) for home care services. Not used for home health services provided by a DME supplier or from a Home IV provider for home IV services.
- 07 = Left against medical advice or discontinued care.
- 09 = Admitted as an Inpatient to this Hospital. Patient admitted to the same short-term medical or specialty hospital where the hospital-based ambulatory surgery service was performed (excluding chronic disease hospitals).
- 20 = Expired.
- 21 = Discharged/transferred to Court/Law Enforcement.
- 50 = Hospice – Home.
- 51 = Hospice – Medical Facility (Certified) Providing Hospice Level of Care.
- 61 = Discharged/transferred to Hospital-Based Medicare Approved Swing Bed.
- 62 = Discharged/transferred to an Inpatient Rehabilitation Facility (IRF), including Rehabilitation Distinct Part Unit of a hospital.

- 63 = Discharged/transferred to a Medicare Certified Long Term Care Hospital (LTCH).
- 64 = Discharged/transferred to a Nursing Facility Certified under Medicaid but not certified under Medicare.
- 65 = Discharged/transferred to a Psychiatric Hospital or Psychiatric Distinct Part Unit of a Hospital.
- 66 = Discharged/transferred to a Critical Access Hospital (CAH).
- 69 = Discharged/transferred to a Designated Disaster Alternative Care Site.
- 70 = Discharged/transferred to another Type of Health Care Institution not defined Elsewhere in this Code List.
- 81 = Discharged to Home or Self Care with a Planned Acute Care Hospital Inpatient Readmission.
- 82 = Discharged/transferred to a Short-Term General Hospital for Inpatient Care with a Planned Acute Care Hospital Inpatient Readmission.
- 83 = Discharged/transferred to Skilled Nursing Facility (SNF) with Medicare Certification with a Planned Acute Care Hospital Inpatient Readmission.
- 84 = Discharged/transferred to a Facility that Provides Custodial or Supportive Care with a Planned Acute Care Hospital Inpatient Readmission.
- 85 = Discharged/transferred to a Designated Cancer Center or Children's Hospital with a Planned Acute Care Hospital Inpatient Readmission.
- 86 = Discharged/transferred to Home under Care of Organized Home Health Service Organization with a Planned Acute Care Hospital Inpatient Readmission.
- 87 = Discharged/transferred to Court/Law Enforcement with a Planned Acute Care Hospital Inpatient Readmission.
- 88 = Discharged/transferred to a Federal Health Care Facility with a Planned Acute Care Hospital Inpatient Readmission.
- 89 = Discharged/transferred to Hospital-Based Medicare Approved Swing Bed with a Planned Acute Care Hospital Inpatient Readmission.
- 90 = Discharged/transferred to an Inpatient Rehabilitation Facility (IRF) including Rehabilitation Distinct Part Units of a Hospital with a Planned Acute Care Hospital Inpatient Readmission.
- 91 = Discharged/transferred to a Medicare Certified Long Term Care Hospital (LTCH) with a Planned Acute Care Hospital Inpatient Readmission.
- 92 = Discharged/transferred to a Nursing Facility Certified under Medicaid but not Certified under Medicare with a Planned Acute Care Hospital Inpatient Readmission.
- 93 = Discharged/transferred to a Psychiatric Hospital or Psychiatric Distinct Part Unit of a Hospital with a Planned Acute Care Hospital Inpatient Readmission.
- 94 = Discharged/transferred to a Critical Access Hospital (CAH) with a Planned Acute Care Hospital Inpatient Readmission.
- 95 = Discharged/transferred to another Type of Health Care Institution not Defined Elsewhere in this Code List with a Planned Acute Care Hospital Inpatient Readmission.

Dataset Segment:**Demographic Variables**

Data Element Name:

Ethnicity

Template Variable:

ethnicity

Format – Length:

Set– maximum 5 codes

Mandatory:

Yes

Description:

Indicates the code that best describes the ethnicity of the patient from the electronic health record (EHR).

Codes and Values:

Examples:

E1 = SPANISH/HISPANIC ORIGIN

E1.04.004 = Colombian

E2 = NOT HISPANIC OR LATINO

E9 = UNKNOWN

Notes for Abstraction:

- If reporting multiple ethnicity codes (up to 5 codes), separate each code using a colon (e.g., “E1.02: E1.04” is Mexican and South American).
- Multiple ethnicity codes within the same heading are expected as there might be many different origins within a heading (e.g., “E1.02.001 Mexican American” and “E1.02.002 Mexicano” are within the same heading “E1.02 Mexican”). However, we would not expect a selection of codes within any two headings of “E1 SPANISH/HISPANIC ORIGIN”, “E2 NOT HISPANIC OR LATINO”, and “E9 UNKNOWN”.
- To obtain the full list of codes, please refer to the following link to the SPARCS code set:
- SPARCS (RR-Race and Ethnicity Codes, Source: Race and Ethnicity Code Set -Version 1.0): <https://www.health.ny.gov/statistics/sparcs/sysdoc/aprr.htm>

Dataset Segment:**Demographic Variables**

Data Element Name:

Facility Identifier

Template Variable:

facility_identifier

Format – Length:

Varchar – 6

Mandatory:

Yes

Description:

This number is the facility's four to six-digit Permanent Facility Identifier (PFI) assigned by the Department of Health.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Enter the Facility Identifier.

Notes for Abstraction:

- Must be a valid number as maintained by the NYSDOH.
- Can only contain numbers 0-9.

Dataset Segment:**Demographic Variables**

Data Element Name:

Gender

Template Variable:

gender

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the gender of the patient.

Codes and Values:

M = Male

F = Female

U = Unknown

Notes for Abstraction:

Dataset Segment:**Demographic Variables**

Data Element Name:	ICD-10-CM Code (n)
Template Variable:	icd_10_cm_code_n
Format – Length:	Set — 8
Mandatory:	Yes

Description:

All diagnosis codes (primary and secondary) from the final hospital billed codes. There can be up to 25 codes, and each code will have its own variable and POA indicator. The first ICD-10-CM (Code 1) will be the **principal** diagnosis.

Codes and Values:

Enter the ICD-10-CM Codes.

Notes for Abstraction:

- These should be reported as 25 individual variables. The variable fields for these will be as follows:
 - The first Data Element will be ICD-10-CM Code 1 with a template variable of icd_10_cm_code1. ICD-10-CM Code 1 is the PRINCIPAL Diagnosis. All other codes will be secondary diagnosis codes.
 - The twentieth Data Element will be ICD-10-CM Code 20 with a template variable of icd_10_cm_code_20.
- Please provide the final hospital billed codes in this field.
- Hospitals may report up to 25 codes and their indicators, including the principal and secondary codes.
- The ICD-10-CM codes would be submitted WITH the appropriate decimal place (AFTER the 3rd character) for each ICD-10-CM code.

Dataset Segment:**Demographic Variables**

Data Element Name:

ICD-10-CM POA Indicator (n)

Template Variable:

icd_10_cm_poa_indicator_n

Format – Length:

Enumerated — 1

Mandatory:

Yes

Description:

Present on Admission (POA) indicator for each ICD-10-CM diagnosis code, aligning with the data element *ICD-10-CM Code (n)*. The first ICD-10-CM POA (Indicator 1) will be the **principal** diagnosis POA indicator.

Codes and Values:

Y = Diagnosis was present at time of inpatient admission

N = Diagnosis was not present at time of inpatient admission

U = Documentation insufficient to determine if the condition was present at the time of inpatient admission

W = Clinically undetermined. Provider unable to clinically determine whether the condition was present at the time of inpatient admission

1 = Unreported/Not used. Exempt from POA reporting.

Notes for Abstraction:

- *ICD-10-CM POA Indicator (n)* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.
- *ICD-10-CM POA Indicator (n)* is not applicable to emergency department (ED) or observation only cases. For ED and observation only cases, report as blank.
- These should be reported as 25 individual variables. The variable fields for these will be as follows:
 - The first Data Element will be ICD-10-CM_POA Indicator 1 with a template variable of icd_10_cm_poa_indicator_1. ICD-10-CM POA Code 1 is the PRINCIPAL Diagnosis POA indicator. All other codes will be secondary diagnosis POA indicators.
 - The twentieth Data Element will be ICD-10-CM POA Indicator 20 with a template variable of icd_10_cm_poa_indicator_20.
- Please provide the final hospital billed codes in this field.
- Hospitals may report up to 25 POA indicators.
- Please provide the final hospital billed code's POA indicator in this field. Please ensure it aligns with *ICD-10-CM Code (n)*.
- Hospitals are required to report a POA indicator for each *ICD-10-CM Code* reported.

- For example, if there are five (5) ICD-10_CM codes reported then five (5) ICD-10-CM POA indicators will be required in the data submission.

Dataset Segment:**Demographic Variables**

Data Element Name:

Inclusion COVID-19

Template Variable:

inclusion_covid_19

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has COVID-19 and/or MIS-C.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: ped_inclusion_definition_table_b_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Demographic Variables**

Data Element Name:

Inclusion Septic Shock

Template Variable:

inclusion_septic_shock

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has septic shock.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: ped_inclusion_definition_table_a_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Demographic Variables**

Data Element Name:

Inclusion Severe Sepsis

Template Variable:

inclusion_severe_sepsis

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has severe sepsis.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: ped_inclusion_definition_table_a_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Demographic Variables**

Data Element Name:	Insurance Number
Template Variable:	insurance_number
Format – Length:	Varchar – 19
Mandatory:	Yes

Description:

Indicates the primary insurance policy identification number for the patient.

Codes and Values:

Enter the Insurance Number.

Notes for Abstraction:

- **Insurance Number** is mandatory.
- Blanks are allowed only
 - If Element Payer is not:
 - Medicare ("C")
 - Medicaid ("D")
 - Insurance Company ("F")
 - Blue Cross ("G")
 - Or, in rare instances when values are truly unattainable from the EHR.
- Must be alphanumeric (0-9) (a-z, A-Z).
- Special characters are invalid entries.

Facilities are directed to enter the following values:

Payer	Type of Number
Blue Cross	Enter the information depending on specific Blue Cross Plan needs and contract requirement.
CHAMPUS	Enter the information depending on CHAMPUS regulations.
Medicaid	Enter Medicaid Client Identification Number (CIN) of the insured or case head Medicaid number shown on the Medicaid Identification Card.
Medicare	Enter the patient's Medicare HIC number as shown on the Health Insurance Card, Certificate of Award, Utilization Notice, Temporary Eligibility Notice, and Hospital Transfer Form or as reported by the Social Security Office.

For all other payer types (commercial insurers, etc.) enter the insured's unique number assigned by the payer.

Dataset Segment:**Demographic Variables**

Data Element Name:

Medical Record Number

Template Variable:

medical_record_number

Format – Length:

Varchar – 17

Mandatory:

Yes

Description:

Indicates the number used by the hospital’s Medical Records Department to identify the patient’s permanent medical record file. This number is not the same as the Patient Control Number.

Codes and Values:

Enter the Medical Record Number.

Notes for Abstraction:

- Must not equal zero or blanks.
- Must be alphanumeric (0-9) (a-z, A-Z).
- Special characters are invalid entries.

Dataset Segment:**Demographic Variables**

Data Element Name:

Other Payer

Template Variable:

other_payer

Format – Length:

Varchar – 50

Mandatory:

Yes

Description:

Indicate the other payers for this hospitalization. This aligns with *Payer* source E and/or I.

Codes and Values:

Enter Other Payer.

Notes for Abstraction:

- If either E or I is reported under *Payer*, then *Other Payer* must be completed. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If multiple other payers are to be reported, each payer will be separated by a colon (:).
- Include a code and a description if a code is captured in your EHR.

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient Control Number

Template Variable:

patient_control_number

Format – Length:

Varchar – 20

Mandatory:

Yes

Description:

Indicates the patient's unique number assigned by the provider to facilitate retrieval of individual financial and clinical records and posting of payment.

Codes and Values:

Enter the Patient Control Number.

Notes for Abstraction:

- Must not equal zero or blanks.
- Must be alphanumeric (0-9) (a-z, A-Z).
- Special characters are invalid entries.

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient City

Template Variable:

pat_addr_city

Format – Length:

Varchar – 30

Mandatory:

Yes

Description:

Indicates the city name of the patient's address.

Codes and Values:

Enter the Patient City Name.

Notes for Abstraction:

- *Patient City* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient County Code

Template Variable:

pat_addr_cnty_cd

Format – Length:

Varchar – 5

Mandatory:

Yes

Description:

Indicates the five-digit Federal Information Processing System (FIPS) county code of the patient's address.

Codes and Values:

Enter the Patient County Code.

Notes for Abstraction:

- *Patient County Code* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.
- Must only consist of numbers (0 – 9).
- FIPS county codes: [state-county code list \(census.gov\)](https://www.census.gov/ipeds/data/enr/states/county_codes.html)

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient Street Address Line 1

Template Variable:

pat_addr_line1

Format – Length:

Varchar – 128

Mandatory:

Yes

Description:

Indicates the first line of patient street address.

Codes and Values:

Enter Patient Street Address Line 1.

Notes for Abstraction:

- *Patient Street Address Line 1* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient Street Address Line 2

Template Variable:

pat_addr_line2

Format – Length:

Varchar – 128

Mandatory:

Yes

Description:

Indicates the second line of patient street address.

Codes and Values:

Enter Patient Street Address Line 2.

Notes for Abstraction:

- *Patient Street Address Line 2* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient State

Template Variable:

pat_addr_st

Format – Length:

Varchar – 2

Mandatory:

Yes

Description:

Indicates the patient's state or province code.

Codes and Values:

Enter Patient State.

Notes for Abstraction:

- *Patient State* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.
- For a complete listing of "State Abbreviations" go to the Official United States Postal Service (USPS) Abbreviations Web site: <https://about.usps.com/who-we-are/postal-history/state-abbreviations.htm>
- Must only consist of letters (a-z, A-Z).

Dataset Segment:**Demographic Variables**

Data Element Name:

Patient Zip Code of Residence

Template Variable:

patient_zip_code_of_residence

Format – Length:

Varchar – 10

Mandatory:

Yes

Description:

Indicates the patient’s 9-digit zip code of residence.

Codes and Values:

Enter the Patient Zip Code of Residence.

Notes for Abstraction:

- *Zip Code of Residence* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.
- Format should be xxxxx-xxxx
- If a hospital does not have the four-digit extension to the zip code, then the five-digit zip code should be reported followed by 0000 in the extension (e.g., 11201-0000).
- Must only consist of numbers (0-9).

Dataset Segment:**Demographic Variables**

Data Element Name:

Payer

Template Variable:

payer

Format – Length:

Set – maximum 3 codes

Mandatory:

Yes

Description:

Indicate the codes that identify the payers for this hospitalization. Provide the primary payer first.

Codes and Values:

A = Self-Pay

B = Workers' Compensation

C = Medicare

D = Medicaid

E = Other Federal Program

F = Insurance Company

G = Blue Cross

H = CHAMPUS

I = Other Non-Federal Program

J = Disability

K = Title V

L = Other/Unknown

Notes for Abstraction:

- Report up to 3 payers.
- If either E or I is reported, then *Other Payer* must be completed.
- Each payer will be separated by a colon (:).
- The PRIMARY payer must be listed first.
 - Example:
 - Workers' Compensation as primary payer and Disability: B:J
 - Blue Cross as primary payer, Insurance Company, Other Federal Program: G:F:E
- If the patient has no insurance prior to admission and is pending Medicaid approval during hospitalization, please report the payer as Medicaid.

Dataset Segment:**Demographic Variables**

Data Element Name:	Race
Template Variable:	race
Format – Length:	Set – maximum 56 codes
Mandatory:	Yes

Description:

Indicates the code that best describes the race of the patient based on the electronic health record.

Codes and Values:

Examples:

R2 = Asian

R2.01 = Asian Indian

R5 = White

Notes for Abstraction:

- If reporting multiple race codes, separate each code using a colon (e.g., “R2.12: R2.01” is Korean and Asian Indian).
- To obtain the full list of codes, please refer to the following link to the SPARCS code set:
- SPARCS (RR-Race and Ethnicity Codes, Source: Race and Ethnicity Code Set -Version 1.0): <https://www.health.ny.gov/statistics/sparcs/sysdoc/aprr.htm>

Dataset Segment:**Demographic Variables**

Data Element Name:

Source of Admission

Template Variable:

source_of_admission

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the code that best describes the patient’s origin before coming to the hospital.

Codes and Values:

- 1 = Non-Health Facility Point of Origin: The patient was admitted to this facility from home or from an assisted living facility.
- 2 = Clinic: The patient was referred to this facility as a transfer from a freestanding or non-freestanding clinic.
- 4 = Transfer from a Hospital (Different Facility): The patient was admitted to this facility as a hospital transfer from an acute care facility where he or she was an inpatient or outpatient.
- 5 = Transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF): The patient was admitted to this facility as a transfer from a SNF or ICF where he/she was a resident.
- 6 = Transfer from Another Health Care Facility: The patient was admitted to this facility as a transfer from another type of health care facility that is not defined elsewhere in this code list.
- 8 = Court/Law Enforcement: The patient was admitted to this facility upon the direction of a court of law or upon the request of a law enforcement agency representative.
- 9 = Information Not Available: The means by which the patient was admitted to this hospital was not known.
- E = Transfer from Ambulatory Surgery Center: The patient was admitted to this facility as a transfer from an ambulatory surgery center.
- F = Transfer from Hospice and is Under a Hospice Plan of Care or Enrolled in a Hospice Program: The patient was admitted to this facility as a transfer from a hospice.

Notes for Abstraction:

- If a patient is moved from one area of the hospital to another (e.g., from the Emergency Department to the ICU), the patient is not considered a transfer. The patient is considered a transfer when the patient is moved between different hospitals with discharge and admission at each location and separate billing from each location.
- Assisted Living is reported as 1, Non-Health Facility Point of Origin.

Dataset Segment:**Demographic Variables**

Data Element Name:

Transferred In

Template Variable:

transferred_in

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates if the patient was received as a transfer from another acute care hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Report “1”, if a patient was transferred in (i.e., received from another acute care hospital).
- Report “0”, if a patient was not transferred in.

Dataset Segment:**Demographic Variables**

Data Element Name:

Transferred Out

Template Variable:

transferred_out

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates if the patient was transferred out to another acute care hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Report “1”, if a patient was transferred out (i.e., transferred/discharged to another acute care hospital).
- Report “0”, if a patient was not transferred out.

Dataset Segment:	Demographic Variables
Data Element Name:	Transfer Facility Identifier Receiving
Template Variable:	transfer_facility_id_receiving
Format – Length:	Varchar – 6
Mandatory:	Yes

Description:

If your hospital received a transfer patient from an acute care hospital, report the hospital PFI from which you received that patient. This is the transferring facility’s four to six-digit Permanent Facility Identifier (PFI) assigned by the Department of Health.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Enter the Transfer Facility Identifier Receiving.

Notes for Abstraction:

- **Transfer Facility Identifier Receiving** is mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- Must be a valid number as maintained by the NYSDOH.
- Must only contain numbers (0-9).
- When receiving a patient from an out-of-state facility, please submit the two-digit state identifier <https://www2.census.gov/geo/docs/reference/state.txt> to represent the transfer facility state. This is **ONLY** to be used when patients are received from an out of state hospital, therefore the code for New York will not be accepted for data submission. For example, a patient received from a Connecticut hospital is submitted with the **transfer_facility_id_receiving** of 09.

To find a hospital PFI, please visit:

https://www.health.ny.gov/statistics/sparcs/reports/compliance/pfi_facilities.htm

Dataset Segment:	Demographic Variables
Data Element Name:	Transfer Facility Identifier Sending
Template Variable:	transfer_facility_id_sending
Format – Length:	Varchar – 6
Mandatory:	Yes

Description:

If your hospital is transferring a patient to another acute care hospital, report the hospital’s PFI to which you are sending the patient. This number is the transfer sending facility’s four to six-digit Permanent Facility Identifier (PFI) assigned by the Department of Health.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Enter the Transfer Facility Identifier Sending.

Notes for Abstraction:

- *Transfer Facility Identifier Sending* is mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- Must be a valid number as maintained by the NYSDOH.
- Must only contain numbers (0-9).
- When transferring a patient to an out-of-state facility, please submit the two-digit state identifier <https://www2.census.gov/geo/docs/reference/state.txt> to represent the transfer facility state. This is ONLY to be used when patients are transferred out of state therefore the code for New York will not be accepted for data submission. For example, a patient transferred to a Connecticut hospital is submitted with the *Transfer Facility Identifier Sending* of 09.

To find a hospital PFI, please visit:

https://www.health.ny.gov/statistics/sparcs/reports/compliance/pfi_facilities.htm

Dataset Segment:**Demographic Variables**

Data Element Name:	Transfer Facility Name Receiving
Template Variable:	transfer_facility_nm_receiving
Format – Length:	Varchar – 50
Mandatory:	Yes

Description:

If your hospital received a patient as a transfer from another acute care hospital, report the hospital name from which you received that patient.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Enter the Transfer Facility Receiving.

Notes for Abstraction:

- *Transfer Facility Name Receiving* is mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- Report when *Transfer Facility Identifier Receiving* is not available.

Dataset Segment:**Demographic Variables**

Data Element Name:	Transfer Facility Name Sending
Template Variable:	transfer_facility_nm_sending
Format – Length:	Varchar – 50
Mandatory:	Yes

Description:

If your hospital is transferring a patient to an acute care hospital, report the hospital’s name to which you are sending the patient.

Department regulations state that services must be reported under the physical location where they are provided. Common ownership of different facilities does not change this requirement.

Codes and Values:

Enter the Transfer Facility Name Sending.

Notes for Abstraction:

- *Transfer Facility Identifier Sending* is mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- Report when *Transfer Facility Identifier Sending* is not available.

Dataset Segment:	Demographic Variables
Data Element Name:	Unique Personal Identifier
Template Variable:	unique_personal_identifier
Format – Length:	Varchar – 10
Mandatory:	Yes

Description:

A composite field comprised of portions of the patient’s last name, first name, and social security number.

Codes and Values:

Included below are the individual components of this data element.

1. **"First 2" and "Last 2" characters of the Patient's Last Name.** The birth name of the patient is preferable if it is available on the facility's information system.
2. **"First 2" characters of the Patient's First Name.**
3. **"Last 4" digits of the Patient's Social Security Number.**

NOTE: This data element is not to be confused with *Patient Control Number*, which provides linkage of all record types containing patient-related data for a specific discharge.

Notes for Abstraction:

First and Last Name Components: Must be **UPPERCASE** alpha characters (A-Z). If the last name is less than 4 characters, the first two and last two characters are used even if some characters are repeated.

- If the first name is only 1-character, repeat the same character to meet the “First 2” character requirement of the Patient’s First Name. For instance, “A” would be reported as “AA”.

Included below are examples of how to report some unusual scenarios: A three-character last name, a two-character last name, a name with junior, a one character first name, a last name with an apostrophe, and a hyphenated last name.

- Joe Tan would be reported as TAANJO
- Bill Su Jr. would be reported as SUSUBI
- E John Smith would be reported as SMTHEE
- Bob O'Brien would be reported as OBENBO
- Sue Jones-Davis would be reported as JOISSU

Social Security Number Component: Must be numeric. If no Social Security Number is available, this sub-field must be zeroes (e.g., TAANJO0000).

Joe Tan with Social Security Number 123-456-7890 would be reported as TAANJO7890

Comorbidity/Risk Factor Variables

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Acute Cardiovascular Conditions (POA)

Template Variable:

acute_cardiovascular_conditions_poa

Format – Length:

Set – maximum of 4 codes

Mandatory:

Yes

Description:

Indicates that the patient had an acute cardiovascular event present on or prior to admission/arrival to the hospital. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

1 = Myocardial infarction

2 = Ischemic stroke/Hemorrhagic stroke/Transient ischemic attack (TIA)

3 = Myocarditis secondary to COVID-19

4 = Other

0 = No acute cardiovascular condition

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- Report all that apply.
- Each value will be separated by a colon (:).

- For example:

- To report multiple elements: 1:2

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_acute_cardiovascular_conditions_code_VerD2.0.csv.

- If the patient has one or more of the ICD-10-CM codes associated with the Codes and Values (1 = Myocardial infarction; 2 = Ischemic stroke/Hemorrhagic stroke/Transient ischemic attack (TIA); 3 = Myocarditis secondary to COVID-19; 4 = Other) for **Acute Cardiovascular Conditions (POA)** in the referenced csv file and has a POA indicator, then:

- Report the respective Codes and Values if the POA indicator is:

- Y = Diagnosis was present at time of inpatient admission
 - U = Documentation insufficient to determine if the condition was present at the time of inpatient admission
 - W = Clinically undetermined; Provider unable to clinically determine whether the condition was present at the time of inpatient admission
 - 1 = Unreported/Not used. Exempt from POA reporting

- Report the Codes and Values as “0 = No Acute Cardiovascular Condition” if the POA indicator is:
 - N = Diagnosis was not present at time of inpatient admission
- If the patient has one or more of the ICD-10-CM codes associated with the Codes and Values (1 = Myocardial infarction; 2 = Ischemic stroke/Hemorrhagic stroke/Transient ischemic attack (TIA); 3 = Myocarditis secondary to COVID-19; 4 = Other) for *Acute Cardiovascular Conditions (POA)* in the referenced csv file and does **not** have a POA indicator, then:
 - Report the respective Codes and Values if the diagnosis was present on admission/arrival, or if documentation is insufficient to determine if the condition was present on admission/arrival, or if it is clinically undetermined whether the condition was present on admission/arrival
 - Report the Codes and Values as “0 = No acute cardiovascular condition” if it is determined that the condition(s) was **not** present on admission/arrival.
- Report the Codes and Values as “0 = No acute cardiovascular condition”, if the patient does not have any of the ICD-10-CM codes associated with the Codes and Values for *Acute Cardiovascular Conditions (POA)* listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Acquired Absent Spleen

Template Variable:

acquired_absent_spleen

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has an acquired absent spleen.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: `ped_acquired_absent_spleen_code_VerD2.0.csv`.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced `csv file`.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced `csv file`.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Adrenal Gland Disorder (POA)

Template Variable:

adrenal_gland_disorder_poa

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has adrenal gland disorder present on or prior to admission/arrival to the hospital. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_adrenal_gland_disorder_code_VerD2.0.csv.
- If the patient has one or more of the ICD-10-CM codes for the **Adrenal Gland Disorder (POA)** listed in the referenced csv file, and has a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the POA indicator is:
 - Y = Diagnosis was present at time of inpatient admission
 - U = Documentation insufficient to determine if the condition was present at the time of inpatient admission
 - W = Clinically undetermined; Provider unable to clinically determine whether the condition was present at the time of inpatient admission
 - Report the Codes and Values as “0 = No”, if the POA indicator is:
 - N = Diagnosis was not present at time of inpatient admission
- If the the patient has one or more of the ICD-10-CM codes for **Adrenal Gland Disorder (POA)** in the referenced csv file and does **not** have a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the diagnosis was present on admission/arrival, or if documentation is insufficient to determine if the condition was present on admission/arrival, or if it is clinically undetermined whether the condition was present on admission/arrival.

- Report the Codes and Values as “0 = No” if it is determined that the condition was **not** present on admission/arrival
- Report the Codes and Values as “0 = No”, if the patient does not have any of the ICD-10-CM codes for *Adrenal Gland Disorder (POA)* listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

AIDS/HIV Disease

Template Variable:

aids_hiv_disease

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has AIDS or an HIV infection.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_aids_hiv_disease_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Asthma

Template Variable:

asthma

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has asthma.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_asthma_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Bronchopulmonary Dysplasia

Template Variable:

bronchopulmonary_dysplasia

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has bronchopulmonary dysplasia (BPD).

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_bronchopulmonary_dysplasia_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Chronic Liver Disease

Template Variable:

chronic_liver_disease

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has chronic liver disease.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_chronic_liver_disease_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Chronic Kidney Disease

Template Variable:

chronic_kidney_disease

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has chronic kidney disease.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_chronic_kidney_disease_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Chronic Respiratory Failure

Template Variable:

chronic_respiratory_failure

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has chronic respiratory failure.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_chronic_respiratory_failure_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Coagulopathy (POA)

Template Variable:

coagulopathy_poa

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has coagulopathy present on or prior to admission/arrival to the hospital. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_coagulopathy_code_VerD2.0.csv.
- If the patient has one or more of the ICD-10-CM codes for the *Coagulopathy (POA)* listed in the referenced csv file, and has a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the POA indicator is:
 - Y = Diagnosis was present at time of inpatient admission
 - U = Documentation insufficient to determine if the condition was present at the time of inpatient admission
 - W = Clinically undetermined; Provider unable to clinically determine whether the condition was present at the time of inpatient admission
 - Report the Codes and Values as “0 = No”, if the POA indicator is:
 - N = Diagnosis was not present at time of inpatient admission
- If the the patient has one or more of the ICD-10-CM codes for *Coagulopathy (POA)* in the referenced csv file and does **not** have a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the diagnosis was present on admission/arrival, or if documentation is insufficient to determine if the condition was present on admission/arrival, or if it is clinically undetermined whether the condition was present on admission/arrival.
 - Report the Codes and Valus as “0 = No” if it is determined that the condition was **not** present on admission/arrival

- Report the Codes and Values as “0 = No”, if the patient does not have any of the ICD-10-CM codes for *Coagulopathy (POA)* listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Congestive Heart Failure

Template Variable:

congestive_heart_failure

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has congestive heart failure.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: `ped_congestive_heart_failure_code_VerD2.0.csv`.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced `csv file`.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced `csv file`.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Cystic Fibrosis

Template Variable:

cystic_fibrosis

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has cystic fibrosis.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_cystic_fibrosis_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Diabetes

Template Variable:

diabetes

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has diabetes.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_diabetes_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Dialysis Comorbidity (POA)

Template Variable:

dialysis_comorbidity_poa

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient arrived at the hospital already receiving dialysis. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_dialysis_comorbidity_code_VerD2.0.csv.
- If the patient has one or more of the ICD-10-CM codes for the *Dialysis Comorbidity (POA)* listed in the referenced csv file, and has a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the POA indicator is:
 - Y = Diagnosis was present at time of inpatient admission
 - U = Documentation insufficient to determine if the condition was present at the time of inpatient admission
 - W = Clinically undetermined; Provider unable to clinically determine whether the condition was present at the time of inpatient admission
 - 1 = Unreported/Not used. Exempt from POA reporting
 - Report the Codes and Values as “0 = No” if the POA indicator is:
 - N = Diagnosis was not present at time of inpatient admission
- If the the patient has one or more of the ICD-10-CM codes for *Dialysis Comorbidity (POA)* in the referenced csv file and does **not** have a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the diagnosis was present on admission/arrival, or if documentation is insufficient to determine if the condition was present on admission/arrival, or if it is clinically undetermined whether the condition was present on admission/arrival.

- Report the Codes and Values as “0 = No” if it is determined that the condition was **not** present on admission/arrival.
- Report the Codes and Values as “0 = No”, if the patient does not have any of the ICD-10-CM codes for *Dialysis Comorbidity (POA)* listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

History of COVID -19

Template Variable:

history_of_covid

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has a history of a positive COVID-19 test within 12 weeks prior to admission/arrival to the hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- Report “1” when the patient has a history of a positive COVID-19 test within 12 weeks prior to arrival at the hospital. If there is a positive test, even if later followed by a negative test, then report the positive test date.
- Report “1” if there is a patient-reported history of COVID-19 within 12 weeks prior to arrival at the hospital, without a supporting LOINC code.
- SARS-Cov-2 LOINC codes can be downloaded to a csv. This file can be found here: <https://loinc.org/sars-cov-2-and-covid-19/>
- These codes are not static and are updated regularly; therefore, hospitals should take care to use the most current list of codes to capture COVID-19 testing.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

History of COVID-19 Datetime

Template Variable:

history_of_covid_dt

Format – Length:

Datetime – 16

Mandatory:

Yes

Description:

Indicates the date and time of the positive *History of COVID-19* test.

Codes and Values:

Enter the History of COVID-19 Datetime.

Notes for Abstraction:

- If there is more than one positive COVID-19 test, report the earliest positive test.
- If there is a patient-reported history of COVID-19 without a supporting LOINC code, report “1” to *History of COVID-19* and leave this variable, *History of COVID-19 Datetime* blank.
- If the *History of COVID-19* is reported as “0 = No”, then *History of COVID-19 Datetime* should be reported as blank.
- *History of COVID-19 Datetime* is mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

History of Other Cardiovascular Disease

Template Variable:

history_of_other_cvd

Format – Length:

Set – maximum 6 codes

Mandatory:

Yes

Description:

Indicates the patient’s history of other cardiovascular disease.

Codes and Values:

1 = Coronary heart disease (e.g., angina pectoris, coronary atherosclerosis)

2 = Peripheral artery disease

3 = Valve disorder

4 = Cerebrovascular disease

5 = Congenital heart defects

6 = Cardiomyopathy

0 = No history of coronary heart disease, peripheral artery disease, valve disorder, cerebrovascular disease, congenital heart defects, or cardiomyopathy.

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2:3
- History of (not acute presentation)
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_history_of_other_cvd_code_VerD2.0.csv.
- Report “0”, if the patient does not have one of the ICD-10-CM code(s) listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Hypertension

Template Variable:

hypertension

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has hypertension.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_hypertension_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Immunocompromising

Template Variable:

immunocompromising

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has an immunocompromising disease/illness.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_immunocompromising_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:	Kawasaki/Mucocutaneous Lymph Node Syndrome
Template Variable:	kawasaki_mucocutaneous
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates that the patient has a diagnosis of Kawasaki/Mucocutaneous lymph node syndrome present on admission/arrival or during hospitalization.

Codes and Values:

0 = No
1 = Yes

Notes for Abstraction:

- POA and/or during hospitalization
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: [ped_kawasaki_mucocutaneous_comorbidity_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced [csv file](#).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced [csv file](#).

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:	Lymphoma Leukemia Multiple Myeloma
Template Variable:	lymphoma_leukemia_multi_myeloma
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates that the patient has malignant neoplasm of lymphatic and hematopoietic tissue including those neoplasms which may be in clinical remission.

Codes and Values:

0 = No
1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_lymphoma_leukemia_multi_myeloma_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Mechanical Ventilation Comorbidity (POA)

Template Variable:

mechanical_vent_comorbidity_poa

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient arrived at the hospital on mechanical ventilation. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_mechanical_vent_comorbidity_code_VerD2.0.csv.
- If the patient has one or more of the ICD-10-CM codes for *Mechanical Ventilation Comorbidity (POA)* listed in the referenced csv file, and has a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the POA indicator is:
 - Y = Diagnosis was present at time of inpatient admission
 - U = Documentation insufficient to determine if the condition was present at the time of inpatient admission
 - W = Clinically undetermined; Provider unable to clinically determine whether the condition was present at the time of inpatient admission
 - 1 = Unreported/Not used. Exempt from POA reporting
 - Report the Codes and Values as “0 = No” if the POA indicator is:
 - N = Diagnosis was not present at time of inpatient admission
- If the the patient has one or more of the ICD-10-CM codes for *Mechanical Ventilation Comorbidity (POA)* in the referenced csv file and does **not** have a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the diagnosis was present on admission/arrival, or if documentation is insufficient to determine if the condition was present on admission/arrival, or if it is clinically undetermined whether the condition was present on admission/arrival.

- Report the Codes and Values as “0 = No” if it is determined that the condition was **not** present on admission/arrival.
- Report the Codes and Values as “0 = No”, if the patient does not have any of the ICD-10-CM codes for *Mechanical Ventilation Comorbidity (POA)* listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Medication Immune Modifying Pre-Hospital

Template Variable:

medication_immune_modifying_pre_hospital

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient is taking disease modifying medications and therapies (drugs and biologics) for collagen diseases, corticosteroids, chemotherapeutic agents through any modality (oral, IV, IM, etc.) known to specifically adversely impact the function of the immune system as the primary therapeutic goal or as an unintended side effect, including steroids (excluding inhaled or topical steroids), and chemotherapy at time of admission.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, medication list, etc.
- For a list of applicable NDC codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: [ped_medication_immune_modifying_ndc_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the NDC codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the NDC codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Metastatic Cancer

Template Variable:

metastatic_cancer

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has metastatic cancer.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_metastatic_cancer_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Neutropenic Patients (POA)

Template Variable:

neutropenic_patients_poa

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has neutropenia present on or prior to admission/arrival to the hospital. This may or may not be evident as a yes on the present on admission indicator as an ICD-10-CM code at or on arrival. Since the intent is to capture all codes not just final billed ICD-10-CMs, a POA indicator may not be available for all cases.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_neutropenia_code_VerD2.0.csv.
- If the patient has one or more of the ICD-10-CM codes for *Neutropenic Patients (POA)* listed in the referenced csv file, and has a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the POA indicator is:
 - Y = Diagnosis was present at time of inpatient admission
 - U = Documentation insufficient to determine if the condition was present at the time of inpatient admission
 - W = Clinically undetermined; Provider unable to clinically determine whether the condition was present at the time of inpatient admission
 - 1 = Unreported/Not used. Exempt from POA reporting
 - Report the Codes and Values as “0 = No” if the POA indicator is:
 - N = Diagnosis was not present at time of inpatient admission
- If the the patient has one or more of the ICD-10-CM codes for *Neutropenic Patients (POA)* in the referenced csv file and does **not** have a POA indicator, then:
 - Report the Codes and Values as “1 = Yes” if the diagnosis was present on admission/arrival, or if documentation is insufficient to determine if the condition was present on admission/arrival, or if it is clinically undetermined whether the condition was present on admission/arrival.
 - Report the Codes and Values as “0 = No” if it is determined that the condition was **not** present on admission/arrival.

- Report the Codes and Values as “0 = No”, if the patient does not have any of the ICD-10-CM codes for *Neutropenic Patients (POA)* listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Obesity

Template Variable:

obesity

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient is obese.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- Combination of ICD-10-CM and/or BMI values from the electronic health record (EHR). Please use the first value upon admission/arrival or the earliest value.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_obesity_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "1", if the patient has a BMI that is in the 95th percentile or higher for age even if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.
- Report "1", if the patient has a BMI value of 30 or higher in the EHR even if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file, does not have a BMI that is in the 95th percentile or higher for age in the EHR, or does not have a BMI value of 30 or higher in the EHR.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Patient Care Considerations

Template Variable:

patient_care_considerations

Format – Length:

Set – maximum 2 codes

Mandatory:

Yes

Description:

Indicates whether the patient has a Do Not Resuscitate (DNR), Do Not Intubate (DNI) or both at any time during the hospital encounter.

Codes and Values:

1 = DNR

2 = DNI

0 = None

Notes for Abstraction:

- This may be present on admission/arrival.
- This may be present at any time during the hospital encounter.
- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Patient Care Considerations Date

Template Variable:

patient_care_considerations_date

Format – Length:

Date – 10

Mandatory:

Yes

Description:

Indicate the earliest date associated with *patient_care_considerations*.

Codes and Values:

Enter the Patient Care Considerations Date.

Notes for Abstraction:

- Format must be YYYY-MM-DD
 - a. YYYY = four-digit year
 - b. MM = two-digit month (01 = January, etc.)
 - c. DD = two-digit day of month (01 through 31)
- Example: November 3, 1959 = 1959-11-03
- If multiple values selected for *patient_care_considerations*, report the earliest date/time associated with the value(s).
- *Patient Care Considerations Date* is mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Pregnancy Comorbidity

Template Variable:

pregnancy_comorbidity

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has a pregnancy-related comorbidity.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- Must be “0” if *Pregnancy Status During Hospitalization* is “0”
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_pregnancy_comorbidity_code_VerD2.0.csv.
- Report “1”, if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report “0”, if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Pregnancy Status During Hospitalization

Template Variable:

pregnancy_status

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient is pregnant, in childbirth, or postpartum on arrival to the hospital or during hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, tests/labs, etc.
- This can be a POA or not a POA variable.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_pregnancy_status_code _VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "1" if detection of human chorionic gonadotropic (hCG) in the urine or blood test (lab value).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file or does not have detection of human chorionic gonadotropic (hCG) in the urine or blood test (lab value).

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:	Prematurity
Template Variable:	prematurity
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates that the patient was born before 37 weeks of pregnancy (preterm birth).

Codes and Values:

- 0 = No
- 1 = Yes
- 2 = NA, Patient > 1-year-old

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_prematurity_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.
- If the patient is greater than 1-year-old, do not specify whether the patient has a history of prematurity. Report these cases as Value "2."

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Sickle Cell Disease

Template Variable:

sickle_cell_disease

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has Sickle Cell Disease.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_sickle_cell_disease_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Skin Disorders/Burns

Template Variable:

skin_disorders_burns

Format – Length:

Set– maximum 3 codes

Mandatory:

Yes

Description:

Indicates that the patient had one or more of the following skin disorders or burns.

Codes and Values:

0 = None

1 = Epidermolysis bullosa

2 = Burn/Corrosion of skin

3 = Frostbite

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_skin_disorders_burn_disease_code_VerD2.0.csv.
- Report “1”, if the patient has one of the ICD-10-CM codes listed in the referenced csv file.
- Report “0”, if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:	Comorbidity/Risk Factor Variables
Data Element Name:	Smoking Vaping
Template Variable:	smoking_vaping
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates that the patient is a current smoker and/or a current vaper.

Codes and Values:

- 0 = No
- 1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_smoking_vaping_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Comorbidity/Risk Factor Variables**

Data Element Name:

Stem Cell Transplant Recipients

Template Variable:

stem_cell_transplant_recipients

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient is a stem cell transplant recipient.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Source is not limited to the billing codes. This information can be obtained from other EHR sources, such as the problem list, medical history, etc.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: ped_stem_cell_transplant_recipients_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Clinical Variables

Dataset Segment:**Clinical Variables**

Data Element Name:

COVID-19 Exposure

Template Variable:

covid_exposure

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient has exposure to COVID-19.

Codes and Values:

0 = No Positive COVID-19 exposure

1 = Positive COVID-19 exposure

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: clinical: ped_covid_exposure_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Clinical Variables**

Data Element Name:

COVID-19 Virus

Template Variable:

covid_virus

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates COVID-19 virus is identified or not identified.

Codes and Values:

0 = COVID-19, virus not identified

1 = COVID-19, virus identified

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: clinical: ped_covid_virus_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Clinical Variables**

Data Element Name:

Drug Resistant Pathogen

Template Variable:

drug_resistant_pathogen

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has resistance to an antimicrobial drug.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: clinical: ped_drug_resistant_pathogen_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Clinical Variables**

Data Element Name:

Flu Positive

Template Variable:

flu_positive

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has a positive flu test present on admission/arrival or during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- POA and/or during hospitalization
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: clinical: ped_flu_positive_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "1", if the patient has a positive influenza virus test (lab value).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file or does not have a positive influenza virus test (lab value).

Dataset Segment:**Clinical Variables**

Data Element Name:

Suspected Source of Infection

Template Variable:

suspected_source_of_infection

Format – Length:

Set – maximum 12 codes

Mandatory:

Yes

Description:

The suspected source of infection.

Codes and Values:

- 1 = Septicemia
- 2 = Bacteremia
- 3 = Fungal infection
- 4 = Peritoneal infection
- 5 = Heart infection
- 6 = Upper respiratory infection
- 7 = Lung infection
- 8 = Central nervous system infection
- 9 = Gastrointestinal infection
- 10 = Genitourinary infection
- 11 = Soft tissue infection
- 12 = Other infection source
- 13 = Unknown

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: clinical: `ped_suspected_source_of_infection_code_VerD2.0.csv`.
- If there is not an identified source of infection as specified in `the referenced csv file`, then report “unknown”.
 - Note that “other infection source” is defined in the ICD-10-CM codes provided in the `csv file`.
- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 2:9:12

Treatment (in hospital) Variables

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

Dialysis Treatment

Template Variable:

dialysis_treatment

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has an order for dialysis during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- For a list of applicable ICD-10-PCS codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: ped_dialysis_treatment_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced csv file.
- Report "1", if the patient has an order for dialysis in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-PCS codes listed in the referenced csv file or does not have an order for dialysis in the EHR.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

During Hospital Immune Modifying Medication

Template Variable:

during_hospital_immune_mod_med

Format – Length:

Number – 1

Mandatory:

Yes

Description:

Indicates that the patient has an order for immune-modifying medication during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable NDC codes, please refer to the corresponding csv file for this version of the data dictionary: comorbidity: [ped_medication_immune_modifying_ndc_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the NDC codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the NDC codes listed in the referenced csv file.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

During Hospital Remdesivir

Template Variable:

during_hospital_remdesivir

Format – Length:

Number – 1

Mandatory:

Yes

Description:

Indicates the patient has an order for remdesivir during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Generic: Remdesivir
- Brand name: Veklury and GS-5734
- For a list of applicable ICD-10-PCS codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: ped_during_hospital_remdesivir_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced csv file.
- Report "1", if the patient has an order for the medication listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-PCS codes or the medication listed in the referenced csv file.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

ECMO

Template Variable:

ecmo

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient has an order for extracorporeal membrane oxygenation (ECMO) during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- For a list of applicable ICD-10-PCS codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: ped_ecmo_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced csv file.
- Report "1", if the patient has an order ECMO in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-PCS codes listed in the referenced csv file or does not have an order for ECMO in the EHR.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

High Flow Nasal Cannula

Template Variable:

high_flow_nasal_cannula

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient has an order for high flow nasal cannula at any time during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Report “1” when the patient has an order for high flow nasal cannula at any time during the hospital encounter.
- Report “0”, if the patient does not have an order for high flow nasal cannula at any time during the hospital encounter.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

IVIG

Template Variable:

ivig

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient has an order for intravenous immunoglobulin (IVIG) during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- For a list of applicable ICD-10-PCS codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: ped_ivig_code_VerD2.0.csv
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced csv file.
- Report "1", if the patient has an order IVIG in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file or does not have an order for IVIG in the EHR.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

Mechanical Ventilation Treatment

Template Variable:

mechanical_vent_treatment

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient has an order for mechanical ventilation at any time during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- For a list of applicable ICD-10-PCS codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: ped_mechanical_vent_treatment_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced csv file.
- Report "1", if the patient has an order for mechanical ventilation in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-PCS codes listed in the referenced csv file or does not have an order for mechanical ventilation in the EHR.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:	Non-Invasive Positive Pressure Ventilation
Template Variable:	non_invasive_pos_pressure_vent
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates the patient has an order for non-invasive-positive pressure ventilation (CPAP, BiPAP) during the hospitalization.

Codes and Values:

0 = No
1 = Yes

Notes for Abstraction:

- Combination of ICD-10-PCS or an order from the electronic health record (EHR).
- For a list of applicable ICD-10-PCS codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: [ped_non_invasive_pos_pressure_vent_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the ICD-10-PCS codes listed in the referenced [csv file](#).
- Report "1", if the patient has an order for non-invasive positive pressure ventilation in the EHR even if they do not have one of the ICD-10-PCS codes listed in the referenced [csv file](#).
- Report "0", if the patient does not have one of the ICD-10-PCS codes listed in the referenced [csv file](#) or does not have an order for non-invasive positive pressure ventilation in the EHR.

Dataset Segment:**Treatment (in hospital) Variables**

Data Element Name:

Vasopressor Administration

Template Variable:

vasopressor_administration

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient has an order for vasopressors during the hospitalization.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable NDC codes, please refer to the corresponding csv file for this version of the data dictionary: treatment: ped_vasopressor_administration_ndc_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the medications or NDC codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the medications or NDC codes listed in the referenced csv file.

Outcome (at discharge) Variables

Dataset Segment:**Outcome (at discharge) Variables**

Data Element Name:

Dialysis Outcome

Template Variable:

dialysis_outcome

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient is discharged on dialysis.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- If there was a patient order to have dialysis at discharge as evidenced by dialysis on the discharge date, report “1.”
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: outcome (at discharge): ped_dialysis_outcome_code_VerD2.0.csv.
- Report “1”, if the patient has the ICD-10-CM code listed in the referenced csv file on the date of discharge.
- Report “0”, if the patient does not have the ICD-10-CM code listed in the referenced referenced csv file on the date of discharge.

Dataset Segment:**Outcome (at discharge) Variables**

Data Element Name:

Mechanical Ventilation Outcome

Template Variable:

mechanical_vent_outcome

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates the patient is discharged on mechanical ventilation.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- If there was a patient order to have mechanical ventilation at discharge as evidenced by mechanical ventilation on the discharge date, report “1”.
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: outcome (at discharge): ped_mechanical_vent_outcome_code_VerD2.0.csv.
- Report "1", if the patient has the ICD-10-CM code listed in the referenced csv file on the date of discharge.
- Report “0”, if the patient does not have the ICD-10-CM code listed in the referenced referenced csv file on the date of discharge.

Dataset Segment:**Outcome (at discharge) Variables**

Data Element Name:

Neurological Outcome

Template Variable:

neurological_outcome

Format – Length:

Set – maximum of 2 codes

Mandatory:

Yes

Description:

Indicates the patient is discharged with neurological outcome(s).

Codes and Values:

0 = No Neurological outcome

1 = Neuropathy

2 = Myopathy

Notes for Abstraction:

- Report all that apply at the date of discharge.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: [outcome \(at discharge\): ped_neurological_outcome_code_VerD2.0.csv](#).

Outcome (in hospital) Variables

Dataset Segment:**Outcome (in hospital) Variables**

Data Element Name:

Cardiovascular Outcomes in Hospital

Template Variable:

cv_outcomes_in_hospital

Format – Length:

Set – maximum of 4 codes

Mandatory:

Yes

Description:

Indicates the patient had one or more of the following cardiovascular outcomes during the hospitalization.

Codes and Values:

0 = None

1 = Acute coronary syndrome

2 = Ischemic stroke

3 = Myocarditis secondary to COVID-19

4 = Cardiomyopathy

Notes for Abstraction:

- Report all that apply.
- Each value will be separated by a colon (:).
- For example:
 - To report multiple elements: 1:2:3
- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: outcome (in hospital): ped_cv_outcomes_in_hospital_code_VerD2.0.csv.

Dataset Segment:	Outcome (in hospital) Variables
Data Element Name:	ICU During Hospitalization
Template Variable:	icu_during_hospitalization
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicate if the patient was admitted to the Intensive Care Unit (NICU; PICU; ICU; MICU; SICU; CCU; Neuro-ICU) during the hospitalization.

Codes and Values:

0 = No
1 = Yes

Notes for Abstraction:

- Report “1”, if the patient was admitted at any time to the ICU during the hospital admission.
- Report “0”, if the patient was not admitted to the ICU during the hospital admission.

Severity Variables

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT 1
Template Variable:	aptt_1
Format – Length:	String – 8
Mandatory:	Yes

Description:

Indicates the first activated partial thromboplastin time (aPTT) level collected after arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT 2
Template Variable:	aptt_2
Format – Length:	String – 8
Mandatory:	Yes

Description:

Indicates the second aPTT value collected after arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT 3
Template Variable:	aptt_3
Format – Length:	String – 8
Mandatory:	Yes

Description:

Indicates the third aPTT level collected after arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT Max
Template Variable:	aptt_max
Format – Length:	String – 8
Mandatory:	Yes

Description:

Indicates the first maximum aPTT value collected after arrival to the hospital.

Codes and Values:

Enter the aPTT levels.

Notes for Abstraction:

- **aPTT 1/2/3/Max** are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital
- Must be reported to one decimal place (example 19.8). For example, 30.7 or 30.0; place hold with 0.
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example, <0.1, should be reported as <0.1.
- If the aPTT level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding the aPTT level results:
 - 30.48 is rounded to 30.5
 - 45.43 is rounded to 45.4
 - 61.75 is rounded to 61.8
 - 55.97 is rounded to 56.0
 - **NOT CORRECT:** 61.75 is truncated to 61.7 (this should be rounded to 61.8)

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT Datetime 1
Template Variable:	aptt_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first aPTT level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT Datetime 2
Template Variable:	aptt_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second aPTT level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT Datetime 3
Template Variable:	aptt_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third aPTT level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	aPTT Datetime Max
Template Variable:	aptt_dt_max
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum aPTT level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the aPTT Datetimes.

Notes for Abstraction:

- **aPTT 1/2/3/Max Datetimes** are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any aPTT is reported then the datetime for the aPTT should be reported. For example, if **aPTT 1** has a value, then **aPTT Datetime 1** should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- This element belongs to Organ Dysfunction Hematologic.

- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	Bilirubin Arrival
Template Variable:	bilirubin_arrival
Format – Length:	String – 6
Mandatory:	Yes

Description:

Indicates the first total bilirubin level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Bilirubin Max
Template Variable:	Bilirubin_max
Format – Length:	String – 6
Mandatory:	Yes

Description:

Indicates the first maximum total bilirubin level collected after arrival to the hospital.

Codes and Values:

Enter the actual total bilirubin level. Convert the units to mg/dL if needed.

Notes for Abstraction:

- **Bilirubin Arrival/Max** are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient’s arrival to the hospital.
- Must be reported to one decimal place (example 2.8).
 - If your EHR allows the capture and the extraction of “<” (less than) or “>” (greater than) sign for this lab, please report “<” (less than) or “>” (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example, <0.1, should be reported as <0.1.
- If the total bilirubin level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding total bilirubin level results:

- 2.51 is rounded to 2.5
- .75 is rounded to .8
- 1.97 is rounded to 2.0
- **NOT CORRECT:** .75 is truncated to .7 (this should be rounded to .8)

Dataset Segment:**Severity Variables**

Data Element Name:	Bilirubin Arrival Datetime
Template Variable:	bilirubin_arrival_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first total bilirubin collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Bilirubin Max Datetime
Template Variable:	bilirubin_max_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum bilirubin level collected after arrival to the hospital.

Codes and Values:

Enter the total Bilirubin Datetimes.

Notes for Abstraction:

- ***Bilirubin Arrival/Max Datetimes*** are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Organ Dysfunction Hepatic is reported then the datetime for Organ Dysfunction Hepatic should be reported. For example, if ***Bilirubin Arrival*** has a value, ***Bilirubin Arrival Datetime*** should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient’s arrival to the hospital.

- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	Creatinine Arrival
Template Variable:	creatinine_arrival
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the first creatinine level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Creatinine Max
Template Variable:	creatinine_max
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the first maximum creatinine level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual creatinine level. Convert the units to mg/dL if needed.

Notes for Abstraction:

- ***Creatinine Arrival/Max*** and corresponding datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient’s arrival to the hospital.
- Must be reported to one decimal place (example 2.8).
 - If your EHR allows the capture and the extraction of “<” (less than) or “>” (greater than) sign for this lab, please report “<” (less than) or “>” (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example, <0.1, should be reported as <0.1.
- If the creatinine level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding creatinine level results:

- 2.81 is rounded to 2.8
- 1.75 is rounded to 1.8
- 1.42 is rounded to 1.4
- 2.97 is rounded to 3.0
- **NOT CORRECT:** 1.75 is truncated to 1.7 (this should be rounded to 1.8)

Dataset Segment:**Severity Variables**

Data Element Name:	Creatinine Arrival Datetime
Template Variable:	creatinine_arrival_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first creatinine level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Creatinine Max Datetime
Template Variable:	creatinine_max_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum creatinine level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the Creatinine Datetimes.

Notes for Abstraction:

- ***Bilirubin Arrival/Creatinine Arrival Datetime*** are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Organ Dysfunction Renal is reported then the datetime for Organ Dysfunction Renal value should be reported. For example, if ***Creatinine Arrival*** has a value, ***Creatinine Renal Arrival Datetime*** should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient’s arrival to the hospital.
- **Formatting:**
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid

2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic First
Template Variable:	diastolic_1
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicates the patient's first diastolic blood pressure collected after arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic Second
Template Variable:	diastolic_2
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicate the patient's second diastolic blood pressure collected after arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic Third
Template Variable:	diastolic_3
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicate the patient's third diastolic blood pressure collected after arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic Min
Template Variable:	diastolic_min
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicates the patient’s first minimum diastolic blood pressure collected after arrival to the hospital.

Codes and Values:

Enter the actual Diastolic Values.

Notes for Abstraction:

- Diastolic values are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Diastolic is reported then the datetime for the Diastolic value should be reported. For example, if *Diastolic Second* has a value, then *Diastolic Second Datetime 2* should not be blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MIN values, report the first one after the patient’s arrival to the hospital.
- If the blood pressure is measured as mean arterial pressure (MAP), report the subsequent blood pressure measured as systolic/diastolic pressures.
- Hospitals may report blood pressure reading obtained by either blood pressure cuff or arterial line.
- Formatting:
 1. Format must be a number up to 3 digits.
 2. Example:
 - a. Diastolic blood pressure 80mm Hg should be reported as 80
 - b. Diastolic blood pressure 112 Hg should be reported as 112

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic First Datetime 1
Template Variable:	diastolic_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first diastolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic Second Datetime 2
Template Variable:	diastolic_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second diastolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic Third Datetime 3
Template Variable:	diastolic_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third diastolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Diastolic Datetime Min
Template Variable:	diastolic_dt_min
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first minimum diastolic blood pressure collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Diastolic Datetimes.

Notes for Abstraction:

- Diastolic Datetimes are mandatory. In rare instances when values are truly unattainable from the EHR report missing values as blank.
- If any Diastolic is reported then the datetime for the Diastolic value should be reported. For example, if *Diastolic Second* has a value, then *Diastolic Second Datetime 2* should not be blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MIN values, report the first one after the patient's arrival to the hospital.
- If the blood pressure is measured as mean arterial pressure (MAP), report the subsequent blood pressure measured as systolic/diastolic pressures.
- Hospitals may report blood pressure reading obtained by either blood pressure cuff or arterial line.
- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	INR 1
Template Variable:	inr_1
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the first INR value collected after the patient’s arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR 2
Template Variable:	inr_2
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the second INR level collected after the patient’s arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR 3
Template Variable:	inr_3
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the third INR level collected after the patient’s arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	INR Max
Template Variable:	inr_max
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the first maximum INR level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the INR level

Notes for Abstraction:

- **INR 1/2/3/Max** and corresponding datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- Must be reported to one decimal place (example 1.2 or 11.5).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example, <0.1, should be reported as <0.1.
- If the INR level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding INR level results:
 - 2.48 is rounded to 2.5
 - 11.75 is rounded to 11.8
 - 2.97 is rounded to 3.0
 - **NOT CORRECT:** 11.75 is truncated to 11.7 (this should be rounded to 11.8)

Dataset Segment:**Severity Variables**

Data Element Name:	INR Datetime 1
Template Variable:	inr_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first INR level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	INR Datetime 2
Template Variable:	inr_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second INR level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	INR Datetime 3
Template Variable:	inr_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third INR collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	INR Datetime Max
Template Variable:	inr_dt_max
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum INR level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the INR Datetimes.

Notes for Abstraction:

- *INR 1/2/3/Max Datetimes* are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any INR is reported then the datetime for the INR value should be reported. For example, if *INR 1* has not value, *INR Datetime 1* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- This element belongs to Organ Dysfunction Hematologic.

- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level 1
Template Variable:	lactate_level_1
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the first lactate level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level 2
Template Variable:	lactate_level_2
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the second lactate level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level 3
Template Variable:	lactate_level_3
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the third lactate level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level Max
Template Variable:	lactate_level_max
Format – Length:	String – 4
Mandatory:	Yes

Description:

Indicates the first maximum lactate level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual lactate level using the mmol/L value. Convert from mg/dL if needed.

Notes for Abstraction:

- **Lactate Level 1/2/3/Max** and corresponding datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- Must be reported to one decimal place (example 5.8).
 - If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value. For example, <0.1, should be reported as <0.1.
- If the lactate level was reported by the lab with more than one decimal place, use the rules of rounding to convert the number to one decimal place.
- Do not just truncate the number in order to convert it to one decimal place.
- Examples of rounding lactate level results:
 - 4.81 is rounded to 4.8
 - 4.85 is rounded to 4.9
 - 4.23 is rounded to 4.2
 - 4.97 is rounded to 5.0
 - **NOT CORRECT:** 4.85 is truncated to 4.8 (this should be rounded to 4.9)

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level Datetime 1
Template Variable:	lactate_level_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first lactate level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level Datetime 2
Template Variable:	lactate_level_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second lactate level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level Datetime 3
Template Variable:	lactate_level_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third lactate level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Lactate Level Datetime Max
Template Variable:	lactate_level_dt_max
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum lactate level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Lactate Datetimes.

Notes for Abstraction:

- *Lactate Level 1/2/3/Max Datetimes* are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Lactate Level is reported then the datetime for the Lactate Level value should be reported. For example, if *Lactate Level 1* has a value, *Lactate Level Datetime 1* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.

- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:

Organ Dysfunction Cardiovascular

Template Variable:

organ_dysfunc_cardiovascular

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has cardiovascular organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: severity: [ped_organ_dysfunc_cardiovascular_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced [csv file](#).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced [csv file](#).

Dataset Segment:**Severity Variables**

Data Element Name:

Organ Dysfunction CNS

Template Variable:

organ_dysfunc_cns

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has central nervous system (CNS) organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: severity: ped_organ_dysfunc_cns_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Severity Variables**

Data Element Name:

Organ Dysfunction Hematologic

Template Variable:

organ_dysfunc_hematologic

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has hematologic organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: severity: [ped_organ_dysfunc_hematologic_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced [csv file](#).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced [csv file](#).

Dataset Segment:**Severity Variables**

Data Element Name:

Organ Dysfunction Hepatic

Template Variable:

organ_dysfunc_hepatic

Format – Length:

Enumerated – 1

Mandatory:

Yes

Description:

Indicates that the patient has hepatic organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No

1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: severity: ped_organ_dysfunc_hepatic_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Severity Variables**

Data Element Name:	Organ Dysfunction Renal
Template Variable:	organ_dysfunc_renal
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates that the patient has renal organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No
1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: severity: ped_organ_dysfunc_renal_code_VerD2.0.csv.
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced csv file.
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.

Dataset Segment:**Severity Variables**

Data Element Name:	Organ Dysfunction Respiratory
Template Variable:	organ_dysfunc_respiratory
Format – Length:	Enumerated – 1
Mandatory:	Yes

Description:

Indicates that the patient has respiratory organ dysfunction after arrival to the hospital.

Codes and Values:

0 = No
1 = Yes

Notes for Abstraction:

- For a list of applicable ICD-10-CM codes, please refer to the corresponding csv file for this version of the data dictionary: severity: [ped_organ_dysfunc_respiratory_code_VerD2.0.csv](#).
- Report "1", if the patient has one or more of the ICD-10-CM codes listed in the referenced [csv file](#).
- Report "0", if the patient does not have one of the ICD-10-CM codes listed in the referenced [csv file](#).

Dataset Segment: **Severity Variables**

Data Element Name:	Platelets 1
Template Variable:	platelets_1
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the first platelet level collected after the patient’s arrival to the hospital.

Dataset Segment: **Severity Variables**

Data Element Name:	Platelets 2
Template Variable:	platelets_2
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the second platelet level collected after the patient’s arrival to the hospital.

Dataset Segment: **Severity Variables**

Data Element Name:	Platelets 3
Template Variable:	platelets_3
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the third platelet level collected after the patient’s arrival to the hospital.

Dataset Segment: **Severity Variables**

Data Element Name:	Platelets Min
Template Variable:	platelets_min
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the first minimum platelet level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual Platelet levels. Convert the units to cells/uL if needed.

Notes for Abstraction:

- *Platelets 1/2/3/Min* and corresponding datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MIN values, please report the first one after the patient's arrival to the hospital.
- If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value.
- This element belongs to Organ Dysfunction Hematologic.

- Formatting:
 1. Format must be a string up to 10-digits long.
 2. Example:
 - a. Platelet 320,000/uL should be reported as 320000
 - b. Platelet 60,000/uL should be reported as 60000

Dataset Segment:**Severity Variables**

Data Element Name:	Platelets Datetime 1
Template Variable:	platelets_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first platelet level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Platelets Datetime 2
Template Variable:	platelets_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second platelet level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Platelets Datetime 3
Template Variable:	platelets_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third platelet level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Platelets Datetime Min
Template Variable:	platelets_dt_min
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first minimum platelet level collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Platelet Datetimes.

Notes for Abstraction:

- *Platelets 1/2/3/Min Datetimes* are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Platelets are reported then the datetime for the Platelets value should be reported. For example, if *Platelets 1* has a value, *Platelets Datetime 1* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MIN values, report the first one after the patient's arrival to the hospital.
- This element belongs to Organ Dysfunction Hematologic.
- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment: **Severity Variables**

Data Element Name:	SIRS Heart Rate 1
Template Variable:	sirs_hearttrate_1
Format – Length:	Enumerated— 3
Mandatory:	Yes

Description:

Indicates the first heart rate value collected after the patient’s arrival to the hospital.

Dataset Segment: **Severity Variables**

Data Element Name:	SIRS Heart Rate 2
Template Variable:	sirs_hearttrate_2
Format – Length:	Enumerated— 3
Mandatory:	Yes

Description:

Indicates the second heart rate value collected after the patient’s arrival to the hospital.

Dataset Segment: **Severity Variables**

Data Element Name:	SIRS Heart Rate 2
Template Variable:	sirs_hearttrate_2
Format – Length:	Enumerated— 3
Mandatory:	Yes

Description:

Indicates the third heart rate value collected after the patient’s arrival to the hospital.

Dataset Segment: **Severity Variables**

Data Element Name:	SIRS Heart Rate Max
Template Variable:	sirs_hearttrate_max
Format – Length:	Enumerated— 3
Mandatory:	Yes

Description:

Indicates the first maximum heart rate value collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual Heart Rates.

Notes for Abstraction:

- Heart Rates are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If *SIRS Heart Rate 2*, *SIRS Heart Rate 3*, and/or *SIRS Heart Rate Max* are collected then these values and their corresponding datetimes should be reported.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- Formatting:
 1. Format must be a number up to 3 digits.
 2. Example:
 - a. Heart rate/Pulse 100 beats per minutes (bpm) should be reported as 100
 - b. Heart rate/Pulse 43 beats per minutes (bpm) should be reported as 43

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Heart Rate Datetime 1
Template Variable:	sirs_hearttrate_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first heart rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Heart Rate Datetime 2
Template Variable:	sirs_hearttrate_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second heart rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Heart Rate Datetime 3
Template Variable:	sirs_hearttrate_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third heart rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Heart Rate Datetime Max
Template Variable:	sirs_hearttrate_dt_max
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum heart rate value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Heart Rate Datetimes.

Notes for Abstraction:

- Heart Rate Datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Heart Rate is reported then the datetime for the Heart Rate value should be reported. For example, if *SIRS Heart Rate 2* has a value, then *SIRS Heart Rate Datetime 2* should not be blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.

- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Leukocyte Arrival
Template Variable:	sirs_leukocyte_arrival
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the first white blood cell (WBC) level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Leukocyte Min
Template Variable:	sirs_leukocyte_min
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the first minimum white blood cell (WBC) level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Leukocyte Max
Template Variable:	sirs_leukocyte_max
Format – Length:	String — 10
Mandatory:	Yes

Description:

Indicates the first maximum white blood cell (WBC) level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual White Blood Cell (WBC) counts. Convert the units to cells/uL if needed.

Notes for Abstraction:

- *SIRS Leukocyte Arrival/Min/Max* and corresponding datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.

- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result of the subsequent lab collected.
- When there are multiple identical MIN or MAX values, report the first one after the patient's arrival to the hospital. If your EHR allows the capture and the extraction of "<" (less than) or ">" (greater than) sign for this lab, please report "<" (less than) or ">" (greater than) sign with the value as well. If your EHR does not capture this, you may simply report the numeric value.
- Formatting:
 1. Format must be a string up to 10-digits.
 2. Example:
 - WBC 100,000/uL should be reported as 100000
 - WBC 11,500/uL should be reported as 11500
 - WBC 4,400/uL should be reported as 4400

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Leukocyte Arrival Datetime
Template Variable:	sirs_leukocyte_arrival_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description

Indicates the date and time of the first white blood cell (WBC) collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Leukocyte Min Datetime
Template Variable:	sirs_leukocyte_min_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description

Indicates the date and time of first minimum white blood cell (WBC) level collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Leukocyte Max Datetime
Template Variable:	sirs_leukocyte_max_dt
Format – Length:	Datetime – 16
Mandatory:	Yes

Description

Indicates the date and time of first maximum white blood cell (WBC) level collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the Leukocyte Datetimes.

Notes for Abstraction:

- *SIRS Leukocyte Arrival/Min/Max Datetimes* are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.

- If any SIRS Leukocyte is reported then the datetime for the SIRS Leukocyte value should be reported. For example, if *SIRS Leukocyte Arrival* has a value, *SIRS Leukocyte Arrival Datetime* should not be blank.
- For all labs, if the initial lab collected was contaminated or determined questionable (e.g., did not result), report the result and the date and time of the subsequent lab collected.
- When there are multiple identical MIN or MAX values, report the first one after the patient's arrival to the hospital.
- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
 MM = two-digit month (01 = January, etc.)
 DD = two-digit day of month (01 through 31)
 hh = two digits of hour (00 through 23) (am/pm NOT allowed)
 mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate 1
Template Variable:	sirs_respiratoryrate_1
Format – Length:	Number — 2
Mandatory:	Yes

Description:

Indicates the first respiratory rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate 2
Template Variable:	sirs_respiratoryrate_2
Format – Length:	Number — 2
Mandatory:	Yes

Description:

Indicates the second respiratory rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate 3
Template Variable:	sirs_respiratoryrate_3
Format – Length:	Number — 2
Mandatory:	Yes

Description:

Indicates the third respiratory rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate Max
Template Variable:	sirs_respiratoryrate_max
Format – Length:	Number — 2
Mandatory:	Yes

Description:

Indicates the first maximum respiratory rate value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the actual Respiratory Rates.

Notes for Abstraction:

- Respiratory Rates are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If *SIRS Respiratory Rate 2*, *SIRS Respiratory Rate 3*, and/or *SIRS Respiratory Rate Max* are collected then these values and their corresponding datetimes should be reported.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- Formatting:
 1. Format must be a number up to 2-digits.
 2. Example:
 - a. Respiratory rate 12 breaths per minutes (bpm) should be reported as 12
 - b. Respiratory rate 9 breaths per minutes (bpm) should be reported as 9

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate Datetime 1
Template Variable:	sirs_respiratoryrate_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first respiratory rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate Datetime 2
Template Variable:	sirs_respiratoryrate_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second respiratory rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate Datetime 3
Template Variable:	sirs_respiratoryrate_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third respiratory rate value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Respiratory Rate Datetime Max
Template Variable:	sirs_respiratoryrate_dt_max
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum respiratory rate value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Respiratory Datetimes.

Notes for Abstraction:

- Respiratory Datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Respiratory Rate is reported then the datetime for the Heart Rate value should be reported. For example, if *SIRS Heart Rate 2* has a value, then *SIRS Heart Rate Datetime 2* should not be blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.

- **Formatting:**
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature 1
Template Variable:	sirs_temperature_1
Format – Length:	Enumerated – 5
Mandatory:	Yes

Description:

Indicates the first temperature value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature 2
Template Variable:	sirs_temperature_2
Format – Length:	Enumerated – 5
Mandatory:	Yes

Description:

Indicates the second temperature value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature 3
Template Variable:	sirs_temperature_3
Format – Length:	Enumerated – 5
Mandatory:	Yes

Description:

Indicates the third temperature value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature Max
Template Variable:	sirs_temperature_max
Format – Length:	Enumerated – 5
Mandatory:	Yes

Description:

Indicates the first maximum temperature value collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual Temperature levels using Fahrenheit. Convert from Celsius if needed.

Notes for Abstraction:

- Temperatures are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If *SIRS Temperature 2*, *SIRS Temperature 3*, and/or *SIRS Temperature Max* are collected then these values and their corresponding datetimes should be reported.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.

- Formatting:
 1. Must be numeric to one decimal place (example 98.8)
 2. Example:
 - a. 100.4°F should be reported as 100.4
 - b. 96°F should be reported as 96.0
 - c. 97.6°F should be reported as 97.6

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature Datetime 1
Template Variable:	sirs_temperature_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first temperature value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature Datetime 2
Template Variable:	sirs_temperature_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second temperature value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature Datetime 3
Template Variable:	sirs_temperature_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third temperature value collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	SIRS Temperature Datetime Max
Template Variable:	sirs_temperature_dt_max
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first maximum temperature value collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Temperature Datetimes.

Notes for Abstraction:

- Temperature Datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Temperature is reported then the datetime for the Temperature value should be reported. For example, if *SIRS Temperature 2* has a value, then *SIRS Temperature Datetime 2* should not be blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MAX values, report the first one after the patient's arrival to the hospital.
- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Dataset Segment:	Severity Variables
Data Element Name:	Systolic First
Template Variable:	systolic_1
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicates the patient’s first systolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Second
Template Variable:	systolic_2
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicate the patient’s second systolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Third
Template Variable:	systolic_3
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicate the patient’s third systolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:	Severity Variables
Data Element Name:	Systolic Min
Template Variable:	systolic_min
Format – Length:	Number – 3
Mandatory:	Yes

Description:

Indicates the patient’s first minimum systolic blood pressure collected after the patient’s arrival to the hospital.

Codes and Values:

Enter the actual Systolic Values.

Notes for Abstraction:

- Systolic values are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Systolic is reported then the datetime for the Systolic value should be reported. For example, if *Systolic Second* has a value, then *Systolic Second Datetime 2* should not be blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MIN values, report the first one after the patient’s arrival to the hospital.
- If the blood pressure is measured as mean arterial pressure (MAP), report the subsequent blood pressure measured as systolic/diastolic pressures.
- Hospitals may report blood pressure reading obtained by either blood pressure cuff or arterial line.
- Formatting:
- Format must be a number up to 3 digits.
 1. Example:
 - a. Systolic blood pressure 80mm Hg should be reported as 80
 - b. Systolic blood pressure 112 Hg should be reported as 112

Dataset Segment:**Severity Variables**

Data Element Name:	Systolic First Datetime 1
Template Variable:	systolic_dt_1
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first systolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Systolic Second Datetime 2
Template Variable:	systolic_dt_2
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the second systolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Systolic Third Datetime 3
Template Variable:	systolic_dt_3
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the third systolic blood pressure collected after the patient’s arrival to the hospital.

Dataset Segment:**Severity Variables**

Data Element Name:	Systolic Datetime Min
Template Variable:	systolic_dt_min
Format – Length:	Datetime – 16
Mandatory:	Yes

Description:

Indicates the date and time of the first minimum systolic blood pressure collected after the patient's arrival to the hospital.

Codes and Values:

Enter the Systolic Datetimes.

Notes for Abstraction:

- Systolic Datetimes are mandatory. In rare instances, when values are truly unattainable from the EHR, report missing values as blank.
- If any Systolic is reported then the datetime for the Systolic value should be reported. For example, if *Systolic Second* has a value, then *Systolic Second Datetime 2* cannot be left blank.
- For all vital signs (i.e., diastolic pressure, systolic pressure, heart rate, respiratory rate, temperature), if the initial vital sign collected was invalid or determined questionable (e.g., instrumentation error), report the value and the date and time of the subsequent vital sign collected.
- When there are multiple identical MIN values, report the first one after the patient's arrival to the hospital.
- If the blood pressure is measured as mean arterial pressure (MAP), report the subsequent blood pressure measured as systolic/diastolic pressures.
- Hospitals may report blood pressure reading obtained by either blood pressure cuff or arterial line.
- Formatting:
 1. Format must be YYYY-MM-DD hh:mm
 - a. YYYY-MM-DDThh:mm is also valid
 2. YYYY = four-digit year
MM = two-digit month (01 = January, etc.)
DD = two-digit day of month (01 through 31)
hh = two digits of hour (00 through 23) (am/pm NOT allowed)
mm = two digits of minute (00 through 59)
 3. Example: 11:42 pm November 3, 1959 = 1959-11-03 23:42
 - a. 1959-11-03T23:42 is also valid
 4. Midnight = 00:00, **NOT** 24:00

Change Log

Version D2.0

- Changes from version D1.1 to D2.0 are highlighted in yellow. Of note, minor grammar edits and font/format changes made for consistency purposes have not been highlighted.
- Added to the “Key points to remember during data extraction” page of the data dictionary:
 - Appendices no longer available as of Data Dictionary Version D3.0.
 - Modified paragraph on inclusion criteria.
- Modified the “Inclusion Definition” page of the data dictionary:
 - Updated inclusion definition for severe sepsis, septic shock, and COVID-19/MIS-C cases based on inpatient, ED, or Observation status.
 - Updated inclusion definition to include newborn patients expired during hospitalization, and severe sepsis/septic shock patients evaluated only in the ED or Observation but never admitted.
 - Removed the following ICD-10-CM codes from Table B: COVID-19 inclusion ICD-10-CM codes:
 - M358 – Other specified systemic involvement of connective tissue
 - M3589 – Other specified systemic involvement of connective tissue
- Added eight (8) new variables to the *Demographic Variables* section of the data dictionary:
 - *Inclusion Septic Shock*
 - *Inclusion COVID-19*
 - *Inclusion Severe Sepsis*
 - *Patient City*
 - *Patient County Code*
 - *Patient Street Address Line 1*
 - *Patient Street Address Line 2*
 - *Patient State*
- Removed the following Note for Abstraction for the *Patient Zip Code of Residence* in the *Demographic Variables* section of the data dictionary
 - If the hospital does not have the patient’s zip code of residence (e.g., foreign resident, homeless), then the five-digit zip code should be reported as 00000-0000.
- Added Notes of Abstraction to *Payer* in the *Demographic Variables* section of the data dictionary:
 - If the patient has no insurance prior to admission and is pending Medicaid approval during hospitalization, please report the payer as Medicaid.
- Added Notes of Abstraction to clarify the role of POA indicator for these following *Comorbidity/Risk Factor Variables*:
 - *Acute Cardiovascular Conditions (POA)*
 - *Adrenal Gland Disorder (POA)*
 - *Coagulopathy (POA)*
 - *Dialysis Comorbidity (POA)*

- *Mechanical Ventilation (POA)*
 - *Neutropenic Patients (POA)*
- Modified variable name and description to the following *Comorbidity/Risk Factor Variables*:
 - *Acquired Absent Spleen*
 - *AIDS/HIV Disease*
 - *Asthma*
 - *Bronchopulmonary Dysplasia*
 - *Chronic Liver Disease*
 - *Chronic Kidney Disease*
 - *Chronic Respiratory Failure*
 - *Congestive Heart Failure*
 - *Cystic Fibrosis*
 - *Diabetes*
 - *History of Other Cardiovascular Disease*
 - *Hypertension*
 - *Immunocompromising*
 - *Lymphoma/Leukemia/Multiple Myeloma*
 - *Medication Immune Modifying Pre-Hospital*
 - *Metastatic Cancer*
 - *Obesity*
 - *Pregnancy Comorbidity*
 - *Prematurity*
 - *Sickle Cell Disease*
 - *Skin Disorders/Burns*
 - *Smoking Vaping*
 - *Stem Cell Transplant Recipients*
- Modified Notes of Abstraction of all variables with references to an appendix.
 - With the removal of the Appendix document, the applicable ICD-10/NDC codes are found in the csv file of the corresponding version of the data dictionary.
- Modified the Codes and Values of *ICD-10-CM POA Indicator (n)* in the *Demographic Variables* section of the data dictionary:
 - Aligned the definition of the Codes and Values to the official indicator definitions (CMS).
 - Replaced “E = Exempt from POA reporting” to “1 = Unreported/Not used. Exempt from POA reporting”
- Added Notes of Abstraction to *ICD-10-CM POA Indicator (n)* in the *Demographic Variables* section of the data dictionary:
 - *ICD-10-CM POA Indicator (n)* is mandatory. In rare instances, when the values are truly unattainable from the EHR, report missing values as blank.
 - *ICD-10-CM POA Indicator (n)* is not applicable to emergency department (ED)/observation only cases. For ED/observation only cases, report as blank.

- Added Notes of Abstraction to **Obesity** in the *Comorbidity/Risk Factor Variables* section of the data dictionary:
 - Report “1”, if the patient has a BMI that is in the 95th percentile or higher for age even if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file.
 - Report “0”, if the patient does not have one of the ICD-10-CM codes listed in the referenced csv file, does not have a BMI that is in the 95th percentile or higher for age in the EHR, or does not have a BMI value of 30 or higher in the EHR.
- Modified variable name and the Description and the Notes of Abstraction of **History of COVID-19 (POA)** in the *Comorbidity/Risk Factor Variables* section of the data dictionary to include “within 12 weeks” timeframe.
- Modified variable name and added Notes of Abstraction to **History of COVID-19 Datetime** in the *Comorbidity/Risk Factor Variables* section of the data dictionary:
 - If the **History of COVID-19** is reported as “0 = No”, then **History of COVID-19 Datetime** should be reported as blank.
- Modified variable name of **Chronic Renal Failure (POA)** in the *Comorbidity/Risk Factor Variables* section of the data dictionary to: **Chronic Kidney Disease**.
- Removed the following Note for Abstraction for the **COVID-19 Virus** in the *Clinical Variables* section of the data dictionary:
 - “This applies to both present on admission/arrival (POA) or acquired during hospitalization.”
- Removed the following *Comorbidity/Risk Factor Variables* from the data dictionary and the csv file:
 - **Altered Mental Status (POA)**
 - **Medication Anticoagulation (POA)**
- Removed the following *Treatment Variables* from the data dictionary and the csv file:
 - **During Hospital Anticoagulation**
- Removed the following *Outcome (at discharge) Variables* from the data dictionary and the csv file:
 - **Cardiovascular Outcomes at Discharge**
- Removed the following *Outcome (in hospital) Variables* from the data dictionary and the csv file:
 - **PE/DVT**
- Revised the name of all pediatric csv files to distinguish from the adult csv files by adding “ped” to the beginning of the csv file name
- Revised the file names for the following *Comorbidity Variables* csv files to correct spelling errors:
 - **Sickle Cell Disease (POA)**
 - sickle_cell_disease_code_VerD2.0.csv changed to ped_sickle_cell_disease_code_VerD2.0.csv
 - **Skin Disorders/Burns (POA)**
 - skin_disorders_burn_disease_code_VerD.X.csv changed to ped_skin_disorders_burn_disease_code_VerD2.0.csv

- Revised *Dialysis Treatment* ICD-10-PCS codes on the csv file: dialysis_treatment_code_VerD2.0.csv
 - Removed the following ICD-10-PCS codes:
 - 5A1D00Z – Performance of Urinary Filtration Single
 - 5A1D60Z – Performance of Urinary Filtration Multiple
 - Added the following ICD-10-PCS code:
 - 3E1M39Z – Peritoneal Dialysis

Version D1.1

- Changes from version D1.0 to D1.1 are highlighted in yellow. Of note, minor grammar edits and font/format changes made for consistency purposes have not been highlighted.
- Modified paragraph on hospital transfer in the *Key point to remember during data extraction* page of the data dictionary.
- Added sentence to the first paragraph of the *Inclusion Definition* page of the data dictionary.
- Added new ICD-10 codes to Severe COVID-19 inclusion ICD-10-CM codes in the *Inclusion Definition* page of the data dictionary
- Added eight (8) new variables to the *Demographic Variables* section of the data dictionary.
- Added one (1) new variable to the *Comorbidity/Risk Factor (POA) Variables* section of the data dictionary.
- Removed “(POA)” from the Dataset Segment “Comorbidity/Risk Factor (POA) Variables.”
- Added “(POA)” to the applicable Data Element Name of the Comorbidity/Risk Factor Variables and updated the Description/Notes of Abstraction of these variables.
- Changed mandatory status from “No” to “Yes” to applicable variables and updated the Description/Notes of abstraction of these variables, allowing for blanks to be submitted for missing data.
- Modified the Format – Length of the following variables:
 - *History of Other Cardiovascular Disease*
 - *Patient Care Considerations*
 - *Cardiovascular Outcomes at Discharge*
 - *Cardiovascular Outcomes in Hospital*
 - *Neurological Outcome*
- Modified the name of the Dataset Segment from “during hospitalization” to “in hospital” for the treatment and outcome variables.
- Modified the Codes and Values of *Skin Disorders/Burns (POA)*
- Updated Notes of Abstraction for clarification on these variables:
 - *Insurance Number*
 - *Patient Zip Code of Residence*
 - *Patient Care Considerations*
 - *Pregnancy Status During Hospitalization*

- *Diastolic First, Diastolic Second, Diastolic Third, Diastolic Min*, and corresponding *Datetimes*
- *Lactate Level 1/2/3/Max*
- *Platelets 1/2/3/Min*
- *SIRS Leukocyte Arrival/Min/Max*
- *Systolic First, Systolic Second, Systolic Third, Systolic Min*, and corresponding *Datetimes*